

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

STIC-ILL

*Adonis only*

**From:** Holleran, Anne  
**Sent:** Friday, December 05, 2003 1:39 PM  
**To:** STIC-ILL  
**Subject:** refs. for 09/780,953

*18-*

Please send copies of the following papers:

1. Gold Molecular Neurobiology (1997) 15(3): 285-306
2. Jain Transplant Intl. (2002) 15(12): 630-634
3. Piek Faseb J. (1999) 13(15): 2105-2124

Anne Holleran  
AU: 1642  
Tel: 308-8892  
RM: 8e03

mailbox: 8e12

# ADONIS - Electronic Journal Services

Requested by

Adonis

Article title FK506 and the role of immunophilins in nerve regeneration

Article identifier 0893764898000462

Authors Gold\_B\_G

Journal title Molecular Neurobiology

ISSN 0893-7648

Publisher Humana Press

Year of publication 1997

Volume 15

Issue 3

Supplement 0

Page range 285-306

Number of pages 22

User name Adonis

Cost centre

PCC \$18.00

Date and time Saturday, December 06, 2003 2:34:10 AM

Copyright © 1991-1999 ADONIS and/or licensors.

The use of this system and its contents is restricted to the terms and conditions laid down in the Journal Delivery and User Agreement. Whilst the information contained on each CD-ROM has been obtained from sources believed to be reliable, no liability shall attach to ADONIS or the publisher in respect of any of its contents or in respect of any use of the system.

# FK506 and the Role of Immunophilins in Nerve Regeneration

**Bruce G. Gold**

*Center for Research on Occupational and Environmental Toxicology, and the Department of Cell and Developmental Biology, Oregon Health Sciences University, Portland, OR 97201*

## Abstract

FK506 is a new FDA-approved immunosuppressant used for prevention of allograft rejection in, for example, liver and kidney transplantations. FK506 is inactive by itself and requires binding to an FK506 binding protein-12 (FKBP-12), or immunophilin, for activation. In this regard, FK506 is analogous to cyclosporin A, which must bind to its immunophilin (cyclophilin A) to display activity. This FK506-FKBP complex inhibits the activity of the serine/threonine protein phosphatase 2B (calcineurin), the basis for the immunosuppressant action of FK506. The discovery that immunophilins are also present in the nervous system introduces a new level of complexity in the regulation of neuronal function. Two important calcineurin targets in brain are the growth-associated protein GAP-43 and nitric oxide (NO) synthase (NOS).

This review focuses on studies showing that systemic administration of FK506 dose-dependently speeds nerve regeneration and functional recovery in rats following a sciatic-nerve crush injury. The effect appears to result from an increased rate of axonal regeneration. The nerve regenerative property of this class of agents is separate from their immunosuppressant action because FK506-related compounds that bind to FKBP-12 but do not inhibit calcineurin are also able to increase nerve regeneration. Thus, FK506's ability to increase nerve regeneration arises via a calcineurin-independent mechanism (i.e., one not involving an increase in GAP-43 phosphorylation). Possible mechanisms of action are discussed in relation to known actions of FKBP: the interaction of FKBP-12 with two  $\text{Ca}^{2+}$  release-channels (the ryanodine and inositol 1,4,5-triphosphate receptors) which is disrupted by FK506, thereby increasing  $\text{Ca}^{2+}$  flux; the type 1 receptor for the transforming growth factor- $\beta$  (TGF- $\beta$ 1), which stimulates nerve growth factor (NGF) synthesis by glial cells, and is a natural ligand for FKBP-12; and the immunophilin FKBP-52/FKBP-59, which has also been identified as a heat-shock protein (HSP-56) and is a component of the nontransformed glucocorticoid receptor.

Taken together, studies of FK506 indicate broad functional roles for the immunophilins in the nervous system. Both calcineurin-dependent (e.g., neuroprotection via reduced NO formation) and calcineurin-independent mechanisms (i.e., nerve regeneration) need to be invoked to explain the many different neuronal effects of FK506. This suggests that multiple immunophilins mediate FK506's neuronal effects. Novel, nonimmunosuppressant ligands for FKBP may represent important new drugs for the treatment of a variety of neurological disorders.

**Index Entries:** Calcineurin; cyclosporin A; FK506; FKBP-12; glucocorticoid receptor; immunophilin; immunosuppressant; nerve regeneration.



## The Current Status of Drugs for Nerve Regeneration

The development of new therapeutics to augment nerve regeneration is an area of intense research activity. Some of the agents that have been experimentally examined are the melanocortins, adrenocorticotrophic hormone (ACTH) and  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) (Bijlsma et al., 1981; 1984; De Koning and Gispén, 1987a; Edwards et al., 1985; Sporel-Özkat et al., 1990; Strand and Kung, 1980; Strand and Smith, 1986), the tri-substituted ACTH analog Org 2766 (Sporel-Özkat et al., 1990; De Koning and Gispén, 1987b), testosterone (Jones, 1993), uridine in experimental diabetic neuropathy (Gallai et al., 1992), gangliosides in diabetic animals (Ekström and Tomlinson, 1990), insulin and insulinlike growth factor (Ekström et al., 1989; Kange et al., 1989; Roth et al., 1995), isaxonine (Sebille et al., 1982), and the long chain fatty alcohol *n*-hexacosanol (Starr et al., 1996). Limited clinical trials have been reported with several of these agents. The most widely lauded of these are the gangliosides. However, in those countries where clinical trials have been undertaken (e.g., Britain, Italy, but not the United States), the use of gangliosides has been discouraged or banned because of the development of Guillain-Barré syndrome in a small percentage of patients (Schonhofer, 1991; Figueras et al., 1992; Landi et al., 1993). Similarly, isaxonine (Nerfactor), was withdrawn from clinical trials because of hepatotoxicity (Letteron et al., 1984). Org 2766 has been tried in a limited number of patients, the results to-date being equivocal (Gorio et al., 1993). One agent currently in clinical use that has been shown to speed functional recovery following a sciatic-nerve crush lesion is nifedipine (Gispén et al., 1991), a dihydropyridine calcium antagonist. However, results obtained from animal studies on its regenerative properties have not been impressive; although regeneration begins slightly earlier, ultimately, there was no difference from untreated ani-

mals in the time needed to achieve full recovery (Angelov et al., 1996). Furthermore, the use of this agent for the treatment of nerve regeneration could be limited by possible untoward cardiovascular effects (Murad, 1990). Thus, the search continues for an agent that can be used to enhance nerve regeneration in humans without adverse side effects.

## FK506: Historical Background

FK506 is a new FDA-approved immunosuppressant macrolide drug isolated from *Streptomyces tsukubaensis* (Kino et al., 1987a,b) and used for organ transplantations (Hoffman et al., 1990; Starzl et al., 1987, 1989). The drug was discovered and isolated in 1984 from a soil sample obtained from Tsukuba, Japan, upon screening microbial fermentation broths using a mixed lymphocyte reaction (Kino and Goto, 1993). The first experimental study of FK506 on cardiac transplantation in rats was reported in 1987 by Ochiai and coworkers (Ochiai et al., 1987). Clinical trials, pioneered by Thomas Starzl's group at the University of Pittsburgh began in February 1989 (Starzl et al., 1987). Over the years, FK506 has been shown to possess two important properties that make it superior to cyclosporin A, currently the most widely employed drug for preventing allograft rejection. First, FK506 is a more potent immunosuppressant (by approx 10X) than cyclosporin A in vitro (Kino et al., 1987a,b; Tocci et al., 1989), in animals (Ochiai et al., 1987; Murase et al., 1987; Ochiai et al., 1988; Sakai et al., 1991; Todo et al., 1987, 1988, 1989), and in humans, leading to fewer instances of rejection and retransplantation (Hoffman et al., 1990; Starzl et al., 1989; McDiarmid et al., 1995). Second, initial reports indicated that its toxicity in humans was far less than that associated with cyclosporin A (Starzl et al., 1989; Shapiro et al., 1990). Whereas the incidence of toxicity has subsequently been found to vary among transplant centers (Klintmalm, 1994; Neuhaus et al., 1994), including the development of moderate-to-severe neurotoxicity (including cortical blindness, tremor, seizures,

and encephalopathy) in anywhere from 3–21% of patients (Lopez et al., 1991; Mueller et al., 1994; Wijdicks et al., 1994; Bronster et al., 1995; Vincenti et al., 1996), this discrepancy has been attributed to the tendency to overdose with FK506 (Fung et al., 1996). Consequently, FK506 is beginning to replace cyclosporin A as the drug of choice in the treatment of allograft rejection.

### FK506: Overview of Neuronal Properties

Whereas its immunosuppressive effects alone make this an important new clinical drug, FK506 has also been found to possess a variety of neuronal properties, including protection against ischemic brain injury (Sharkey and Butcher, 1994; Ide et al., 1996; Tokime et al., 1996; Butcher et al., 1997) and glutamate neurotoxicity in vitro (Dawson et al., 1993) (although a very recent study [Butcher et al., 1997] did not substantiate protection against glutamate toxicity in vivo), prevention of N-methyl-D-aspartate (NMDA)-receptor desensitization (Tong et al., 1995), prevention of kindling (Moriwake et al., 1996) blockade of long-term potentiation (LTP) and long-term depression (LTD) in the visual cortex (Torii et al., 1995; Funauchi et al., 1994), facilitation of LTP (Ikegami et al., 1996) and blockade of LTD in the rat hippocampus (Hodgkiss and Kelly, 1995), and alteration in neurotransmitter release (Steiner et al., 1996) and endocytosis (Kuromi et al., 1997). In regard to the latter, whereas LTP and LTD have not been proven to underlie learning and memory; it may be relevant that cyclosporin A also inhibits memory formation in day-old chicks (Bennett et al., 1996). It is likely that all these FK506 neuronal properties are mediated by calcineurin inhibition. For example, calcineurin is known to regulate NMDA-receptor desensitization (see Tong et al., 1995). Furthermore, by preventing calcineurin-dependent NOS dephosphorylation, FK506 would inhibit NOS activity

thereby reducing formation of NO that has been implicated in mediating neurotoxicity (Dawson et al., 1993; Snyder and Sabatini, 1995), and the generation of both LTP (Schuman and Madison, 1991; O'Dell et al., 1991; Kantor et al., 1996) and LTD (Shibuki and Okada, 1996; Malen and Chapman, 1997). The apparent contradictory findings that FK506 elicits both an inhibition of NMDA-induced neurotransmitter release (Steiner et al., 1996) and an augmentation of depolarization-induced neurotransmitter release have been attributed to a similar underlying mechanism; i.e., altered phosphorylation of the calcineurin substrates NOS and synapsin I, respectively (Steiner et al., 1996). It is possible that any one of these alterations may play a role in the development of some of the neurological sequelae observed in the occasional patient undergoing FK506 therapy (Lopez et al., 1991; Mueller et al., 1994; Wijdicks et al., 1994; Bronster et al., 1995; Vincenti et al., 1996).

In addition to these interesting neuronal properties, FK506 speeds nerve regeneration in the peripheral nervous system of the rat with a focal crush lesion of the sciatic nerve (Gold et al., 1994, 1995). The nerve regenerative property of FK506 is the focus of this review. *A priori*, it would not be expected that an immunosuppressant drug (FK506) would alter axonal regeneration since Wallerian degeneration is not an immune-mediated event (Griffin et al., 1993). Furthermore, even if a reduction in macrophage infiltration (Brück and Friede, 1990) occurs following FK506 administration, such an alteration would be expected to impair nerve regeneration by delaying the removal of products of Wallerian degeneration from the distal stump (Beuche and Friede, 1984; Friede and Brück, 1993) studies utilizing the Ola (Wld) mouse mutant (Perry et al., 1990a,b), for example, show that a delay in Wallerian degeneration and macrophage infiltration leads to an impairment in axonal regeneration of sensory (Ludwin and Bisby, 1992) and motor (Chen and Bisby, 1992) neurons. Why, therefore, should FK506 alter nerve regeneration? To answer this question, it is first

necessary to understand FK506's mechanism of immunosuppression.

## The Immunophilins and Immunosuppression

Immunosuppressant drugs FK506 and cyclosporin A are prodrugs that are activated when bound to their respective binding proteins (immunophilins), FK506-binding-protein (FKBP) and cyclophilin, respectively (Schreiber and Crabtree, 1992). The function of the immunophilins as mediators of immunosuppressant drugs has recently been reviewed (Schreiber, 1991). FKBP's are a family of immunophilin proteins named according to their size (in kD): for example, FKBP-12, -13, -25, -51, -52, and -59 (Schreiber and Crabtree, 1992; Alnemri et al., 1993; Sigal and Dumont, 1992). Initially, it was thought that the immunosuppressant actions of FK506 and cyclosporin A were caused by the ability of their respective immunophilins to prevent the interconversion of the *cis*- and *trans*- isomers of prolyl residues of proteins (Walsh et al., 1992). However, subsequent studies proved that peptidyl prolyl isomerase (PPI) activity does not play a role in immunosuppression (Dumont et al., 1992; Wiederrecht et al., 1992), since not all immunosuppressant analogs of these two agents inhibit isomerase activity (Sigal et al., 1991) and, conversely, not all isomerase inhibitors are immunosuppressants (Tocci et al., 1989; Dumont et al., 1992; Sigal et al., 1991; Bierer et al., 1990; for review, see Wiederrecht and Etzkorn, 1995). [The rotamase activity of the immunophilins has been shown to have physiological functions, such as inhibition of collagen assembly by cyclosporin A (Lewin, 1995; Compton et al., 1992; Bächinger et al., 1993) and FK506 (Hans Peter Bächinger, personal communication).] It is widely accepted that the most likely mechanism for the action of these agents involves inhibition of the activity of *calcineurin* (Freman et al., 1992), a calcium/calmodulin-dependent phosphoserine/phosphothreonine protein phosphatase,

also known as PP-2B (Walsh et al., 1992; Dumont et al., 1992; Liu et al., 1991). By inhibiting the calcineurin-induced dephosphorylation of a nuclear factor of activated T-cells (NFAT), FK506 and cyclosporin A prevent NFAT translocation into the nucleus where it induces interleukin-2 (IL-2) secretion, thereby preventing T-cell proliferation (Tocci et al., 1989; Schreiber and Crabtree, 1992; Sigal and Dumont, 1992; Freman et al., 1992; Terada et al., 1992). Of the variety of FKBP's, it has been demonstrated (Sigal and Dumont, 1992; Liu et al., 1991) that FKBP-12 mediates FK506's immunosuppressant activity in T-cells. Whether the inhibition of the expression of the proto-oncogenes *c-myc* and *c-fos* (Sigal and Dumont, 1992) is involved in their immunosuppressant action is unclear. Recently, a second FKBP (FKBP-51), which is specifically expressed in T-lymphocytes (T-cells), has been demonstrated to also inhibit calcineurin (Baughman et al., 1995) suggesting that multiple immunophilins may participate in mediating FK506's immunosuppressant action.

## The Immunophilins in the Nervous System

Immunophilins are enriched in neurons throughout the central (Steiner et al., 1992) and peripheral (Lyons et al., 1995) nervous systems. Their distributions and putative actions in regulating neuronal function have also been reviewed (see Snyder and Sabatini, 1995; Wiederrecht and Etzkorn, 1995; Sánchez and Ning, 1996). Most studied is FKBP-12 (Harding et al., 1989; Siekierka et al., 1989), a ubiquitous protein that has been highly conserved throughout evolution (Siekierka et al., 1990). Snyder and coworkers, who first reported that FKBP-12 is present in neuronal tissue, showed that the mRNA levels for FKBP-12 increase in motor (facial) neurons following peripheral nerve axotomy (Lyons et al., 1992). In addition, calcineurin is present not only in T-cells but also in brain (Steiner

et al., 1992) where it comprises up to 1% of total protein in some brain regions (Klee, 1991). Furthermore, the high levels (relative to other tissues) of this protein in brain (Mukai et al., 1993) correspond regionally with the presence of high levels of FKBP (Steiner et al., 1992). Interestingly, one of the major targets for calcineurin in neurons is the growth-associated protein GAP-43 (Skene, 1990). Both GAP-43 (Skene, 1989; Skene and Willard, 1981a,b) and neuronal phosphatases (Bixby and Jhabvala, 1993) are concentrated in growth cones where they are believed to play an important role in nerve regeneration. Most importantly, a preliminary report (Steiner et al., 1991) indicates that FK506 increases phosphorylation of GAP-43. Based upon this information, FK506 (and, if this model is correct, cyclosporin A) should alter nerve regeneration, perhaps via phosphorylation-dependent activation of GAP-43 (Liu and Storm, 1990). Although this hypothesis initially appeared tenable, the most recent findings from my laboratory (summarized in the next three sections) do not support a calcineurin-dependent mechanism for how FK506 increases the rate of axonal regeneration.

### **FK506 Increases the Rate of Axonal Regeneration in the Rat Sciatic Nerve**

We reported preliminarily in 1993 (Gold et al., 1993) and formally in 1994 (Gold et al., 1994), that FK506 increases functional recovery and nerve regeneration in young adult Sprague-Dawley rats (160–225 g) given a focal sciatic nerve crush (axotomy). A second paper, in 1995, showed that FK506 increases the rate of axonal regeneration in this model (Gold et al., 1995). Axotomy was performed by crushing the nerve twice (for a total of 30 s using a No. 7 Dumont jeweler's forceps) at the level of the hip; the crush site was marked by a sterile 9-0 suture inserted in the epineurial sheath (Gold et al., 1994; 1995). Daily subcutaneous injections of FK506 (1 mg/kg) reduced (by approx 1.5 d) the

number of days until the onset of an ability to right the foot and move the toes ("onset"), and the number of days until the animal demonstrated an ability to walk on its hind feet and toes ("walking") (Fig. 1); animals were observed by at least two independent investigators in a double-blinded study (i.e., the treatment regimen was unknown to both the presenter and the observers of the animals). Control (saline- or vehicle-treated) animals first demonstrated an ability to right the foot between 17 and 18 d. In contrast, three of the five FK506-treated animals walked almost normally by 18 days. These differences in functional recovery were reflected in the morphological appearance of the sciatic nerve and its branches distal to the nerve crush at 18 d after axotomy. Axonal calibers (determined by electron microscopy in plastic-embedded sections from glutaraldehyde-perfused animals) for the largest 30% of regenerating axons were increased by 93% in the tibial nerve branch to the soleus muscle from FK506-treated animals (see Fig. 3 in Gold et al., 1994). Axons were found to have advanced further toward their targets and reinnervation of intrafusal fibers was evident in the most distal (interosseus) muscles. The maximal distance of axonal elongation from the crush site was measured at 10 and 15 d following axotomy using radiolabeling techniques (Gold et al., 1995), the rate of regeneration being estimated from the slope of the resultant line determined between these two-time points. We found that a 1 mg/kg daily dosage of FK506 significantly ( $p < 0.05$ ) increased (by 16%) the axonal regeneration rate from 3.8 mm/d in saline-treated rats to 4.4 mm/d in FK506-treated animals (see Fig. 7 in Gold et al., 1994); regeneration rate was determined using radiolabeling techniques that provide a more accurate and less biased assessment of regenerative distance than the more commonly used pinch test. Taken together, these initial studies demonstrated that FK506 speeds functional recovery by increasing the rate of axonal regeneration in the rat sciatic nerve.

We have recently extended these studies by examining the dose-dependency for FK506's




Fig. 1. Bar graphs showing number of days from axotomy until onset of toe movement and an ability to right the foot (left), and an ability to walk on the hind feet (right) are present in saline-treated and FK506-treated rats. Both signs of functional recovery appear earlier in all FK506-treated groups compared to saline-treated controls, being present earliest in the animals given the 5 mg/kg dose. \* $p < 0.0001$ , compared to saline-treated controls (by one-way ANOVA and Fisher's post-hoc test); † $p < 0.01$ , compared to 1 and 2 mg/kg groups (by one-way ANOVA and Fisher's post-hoc test); †† $p < 0.001$ , compared to 1 and 2 mg/kg groups (by one-way ANOVA and Fisher's post-hoc test).

nerve regenerative effect (Wang et al., 1997). For these studies, we employed daily subcutaneous injections of FK506 at dosages of 2, 5, or 10 mg/kg. In all our analyses, the best results were obtained in animals given the 5 mg/kg dose. In terms of functional recovery, the 5 mg/kg FK506-treated group demonstrated the earliest recovery of function in the hindfeet (Fig. 1); for example, the number of days until the onset of an ability to right the foot and move the toes ("onset"), and the number of days until the animal demonstrated an ability to walk on its hind feet and toes ("walking") were reduced from  $16.8 \pm 0.20$  d to  $14 \pm 0$  d and  $17.8 \pm 0.2$  d to  $15 \pm 0$  d, respectively, in the saline-treated animals ( $n = 5$ ) and in the




Fig. 2. Representative footprints at 18 d following axotomy from a saline-treated rat (A) and an animal given 5 mg/kg FK506 (B). Each image was generated by scanning the original footprint using MacImage (Xerox Imaging Systems, Stamford, CT). The foot and all toes (numbered) are clumped together in the footprint from the saline-treated rat. In contrast, the footprint from the FK506-treated animal exhibits toe spread for all five digits; the lack of an imprint by the heel shows that the animal was able to support its weight on its toes and the front of its foot during walking.

5 mg/kg group ( $n = 3$ ), respectively. Representative footprints at 18 d following axotomy (corresponding to the time of morphological analysis; see below) are shown in Fig. 2. Whereas the footprint from the saline-treated, axotomized control animal shows a continued deficit (Fig. 2A), the footprint from the animal given the 5 mg/kg dose (Fig. 2B) appears virtually normal; this rat demonstrates an ability to walk on toes and the front of its foot, as shown by the lack of a heel imprint. Electron microscopy, performed at 18 d, confirmed the behavioral findings. Nerves from FK506-treated animals contained larger, more advanced regenerating axons, representing the morphological correlate of the earlier functional recovery in these animals. The percent increase in mean axonal areas from control values demonstrated a bell-shaped dose-dependency (Fig. 3); mean axonal areas in FK506-treated animals (including our earlier data from rats treated with 1 mg/kg/d) were increased by 67, 83, 120, and 100% in the 1, 2, 5, and 10 mg/kg groups, respectively. Measurement of axonal regenera-




Fig. 3. FK506 dose-dependently increases mean axonal area in the soleus nerve. The percent increase in axonal areas from control values increases with dose between 1, 2, and 5 mg/kg, declining slightly in the 10 mg/kg group.




Fig. 4. FK506 dose-dependently increases the rate of axonal regeneration in the sciatic nerve. The percent increase in regeneration rate from control values increases with dose between 1 and 5 mg/kg, declining slightly in the 10 mg/kg group.

tion using radiolabeling techniques also exhibited a bell-shaped dose-response (Fig. 4); the percent increase in rate of axonal regeneration from control values was 16, 34, and 29% in the

1, 5, and 10 mg/kg groups, respectively. Taken together, these data establish the dose-dependency for the ability of FK506 to increase nerve regeneration in vivo.

## Cyclosporin A Does Not Increase Axonal Regeneration Rate

To determine if calcineurin is involved in the ability of FK506 to increase the rate of axonal regeneration, we tested cyclosporin A to see if the drug shares FK506's regenerative properties (Wang et al., 1997). Since the drug-immunophilin complexes inhibit calcineurin activity to suppress T-cell function, it would be expected that equivalent doses of cyclosporin A would also increase the rate of axonal regeneration in a dose-dependent manner; i.e., if calcineurin inhibition is the underlying mechanism for FK506's nerve regenerative effect. Dosages (10 and 50 mg/kg/d) of cyclosporin A chosen for study were based upon its relative potency for immunosuppression and PPI (with cyclosporin A being approx 1/10 as potent as FK506) and our finding that a daily dose of 5 mg/kg of FK506 maximally increases the axon regeneration rate; the 50 mg/kg/d dosage approaches the maximal tolerated dose in rats and is sufficient to prevent allograft rejection in rodents (Cosenza et al., 1994). Neither dose of cyclosporin A altered the rate of axonal regeneration (as determined by radiolabeling techniques) nor did the drug speed functional recovery in the sciatic nerve. These findings indicate that cyclophilin A-mediated calcineurin inhibition does not mediate the ability of FK506 to increase nerve regeneration.

## FKBP-12 Ligands Not Inhibiting Calcineurin Increase Nerve Regeneration

To determine whether FK506-FKBP-12 ligands increase nerve regeneration via a calcineurin-independent mechanism, studies have recently been conducted to test the regenerative potential of potent FKBP-12 inhibitors that do not inhibit calcineurin. Snyder and coworkers (Steiner et al., 1997) reported that *topical* administration of L-685,818 (18-OH, 21-ethyl-FK506) to the nerve-crush site accelerates functional

recovery and increases axonal calibers (using paraffin-embedded nerves fixed by immersion in 10% formalin) distal to the crush lesion. Unfortunately, any correlation between functional recovery and morphological changes in these animals is problematic since the investigators limited their assessment of the axons to a distance of only 2 mm from the crush injury at 18 d following the lesion. It is therefore unclear whether the presence of more myelinated fibers is because of an effect of the drug on axonal sprouting as opposed to axonal elongation *per se*, an interpretation supported by their *in vitro* data (Steiner et al., 1997). Furthermore, the recent finding that L-685,818-FKBP-12 is able to inhibit calcineurin in *C. neoformans* (Odom et al., 1997) makes this compound of questionable value for definitively ruling out calcineurin activity in the nervous system.

Based upon the structural domains of FK506 (Fig. 5), two groups have synthesized novel small molecules that retain the FKBP-12 binding domain but *lack* the structural components of the effector domain. Snyder and coworkers reported (Steiner et al., 1997a,b) that *systemic* administration (i.e., subcutaneous injections for 18 d) of two FKBP-12 ligands (with binding affinities of 25- and 250-nM, respectively) but that do not inhibit calcineurin activity (and are not immunosuppressants) increase the size of myelinated fibers (at 2 mm distal to the crush site). As in the case of the L-685,818 compound, it is difficult to rule out an effect of the compounds on axonal sprouting, a possibility apparently supported by their findings in the striatum from *N*-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-treated mice (Steiner et al., 1997b) (*see* FK506: Regeneration of CNS Neurons). This concern is exacerbated by their failure to assess functional recovery in animals given these compounds. These investigators also report an increase in myelination that may merely reflect the increase in axonal calibers, as opposed to a direct effect on myelination. Morphometric analysis of glutaraldehyde-fixed tissue is needed to differentiate between these two possibilities (e.g., calculating the G-ratio: diameter of the axon/diameter of the nerve fiber).




Fig. 5. Comparison of the structures of FK506 (left) and V-10,367 (right). The bracketed portion of FK506 represents the calcineurin-binding domain which is absent in V-10,367.

We have tested (Gold et al., 1997) by *systemic* administration (i.e., subcutaneous injections for 18 d), the nerve-regenerative property of V-10,367 (Fig. 5), a small molecule that also lacks the structural components of the effector domain necessary for calcineurin binding but that binds to FKBP-12 with much greater (<1 nM) affinity (Armistead et al., 1995) than the compounds used by Snyder and coworkers (Steiner et al., 1997a,b). As expected, this compound does not inhibit calcineurin activity (D. R. Armistead, personal communication). Rats given subcutaneous injections of V-10,367 (400 mg/kg/d) showed a more rapid (by approx 2 d) functional recovery following a focal sciatic nerve crush compared to vehicle-treated control animals. The morphological correlate of this earlier return of function was the presence of larger-sized regenerating axons in the sciatic nerve. By electron microscopy, mean axonal areas in the soleus nerve were 50% larger in the V-10,367-treated animals compared to control values. We found no evidence for an increase in myelination, as determined from the G-ratio (B.G. Gold and M.-S. Wang, unpublished obser-

vation). Whereas the increased size of regenerating axons was somewhat less than with FK506, further studies are needed to determine the optimal dosage of this compound. The compound is also effective by oral administration at dosages as low as 5 mg/kg (B. G. Gold, M. Zeleny-Pooley, and M.-S. Wang, unpublished observation). Thus, our study is the first to demonstrate that systemic administration of a potent FKBP-12 ligand that lacks the structural components necessary for calcineurin inhibition speeds functional recovery by accelerating the growth of regenerating axons to the distal musculature following a sciatic nerve-crush lesion. Future studies should address whether the compound also accelerates nerve regeneration in older animals as well as in other species.

### FK506: Effects on Neurite Outgrowth in Culture

In the seminal 1994 study, Snyder and coworkers (Lyons et al., 1994) showed that FK506 potently (as low as 0.1 nM) increases



neuritic outgrowth from PC12 cells and DRG explant cultures in a concentration-dependent fashion. In PC12 cells, neurite outgrowth was assessed by determining the percentage of cells with processes greater than 5  $\mu\text{m}$ . The ability of FK506 to enhance neurite outgrowth in PC12 cells was dependent upon the concentration of NGF. Maximal efficacy was found in the presence of relatively low (1–10 ng/mL) concentrations of NGF, since at higher concentrations of NGF (>10 ng/mL) the maximal response (in terms of % of cells bearing processes) was already demonstrated. Half-maximal response was obtained at 0.1 nM in the presence of submaximal (1 ng/mL) concentrations of NGF. The finding that similar results were obtained in DRG cultures indicates that FK506 activity is not dependent upon exogenously supplied neurotrophins (e.g., NGF) (*see Putative Mechanisms for FK506's Ability to Increase Nerve Regeneration*).

Recently, these investigators reported that cyclosporin A also increases neurite outgrowth in these two systems. We have used human neuroblastoma SH-SY5Y cells to study the concentration-dependency of FKBP-12 ligands for increasing neurite outgrowth. We measured neurite process length at 96 and 168 h after addition of the test agent. Maximal efficacy was observed between 1 and 10 nM in the presence of 10 ng/mL NGF. In contrast to Snyder and coworkers (Steiner et al., 1997a), and in accordance with our *in vivo* data (*see Cyclosporin A Does Not Increase Axonal Regeneration Rate*), we found that cyclosporin A (1–1000 nM) did not significantly alter neurite outgrowth in SH-SY5Y cells (B. G. Gold and M. Zeleny-Pooley, unpublished observations).

The results of these studies stand in contrast to those reported by Jay and coworkers (Chang et al., 1995) who found that FK506 *reduces* neurite outgrowth in cell culture. These studies employed a much higher (50  $\mu\text{M}$ ) concentration of FK506 in the presence of very high levels of NGF (100 ng/mL). In this context, we have found that 1  $\mu\text{M}$  FK506 actually inhibits NGF-induced neurite outgrowth in SH-SY5Y cells (B. G. Gold and

M. Zeleny-Pooley, unpublished results). Thus, in the context of the broad range of concentrations of FK506 we have examined, our results are, in fact, consistent with their findings.

## FK506: Regeneration of CNS Neurons

Cell culture systems provide a simple means to examine whether FK506 also increases nerve regeneration in the CNS. We have conducted *in vitro* studies to examine whether FK506 also promotes axonal regeneration in cells not derived from the PNS. Our studies indicate that FK506 (1–100 nM), in the absence of exogenously supplied neurotrophins (e.g., neurotrophin-3; NT-3) also increased neuritic outgrowth in rat cortical neurons and hippocampal neurons (Fig. 6) in a concentration-dependent manner (B. G. Gold and M. Zeleny-Pooley, unpublished results).

An important question is whether FK506 is also effective *in vivo* in aiding nerve regeneration in the CNS. One preliminary report has examined the ability of daily subcutaneous injections of FK506 (0.5 mg/kg) to increase regeneration of dorsal-root axons into the spinal cord (Sugawara et al., 1995). This study was restricted to the examination of unmyelinated axons immunoreactive for calcitonin gene-related peptide (CGRP). Whether these axons originated in the DRG or represented abnormal sprouting of axons intrinsic to the spinal cord is unclear (Goldberger et al., 1993). Moreover, it is possible that in this model FK506 may have inhibited an autoimmune response leading to increased regeneration via its immunosuppression action.

A recent report (Steiner et al., 1997b) indicates that a nonimmunosuppressant FKBP-12 ligand also enhances the density of striatal dopaminergic innervation (as determined by immunocytochemical staining for tyrosine hydroxylase) in MPTP-intoxicated mice. Importantly, the FKBP-12 ligand increased the density of staining when given after the toxic agent,




Fig. 6. Hippocampal neurons 86 h in culture either untreated (A) or treated with 100 nM FK506 (B). The axonal processes (\*) are clearly discernible and are markedly elongated in the neuron given FK506; neuritic length was 82 and 166  $\mu\text{m}$  in (A) and (B), respectively. Neurons were grown on coverslips that were inverted onto 24-well plates precoated with a monolayer of cortical astrocytes, according to Banker and Cowan (1977). Magnification 680X.

suggesting that the compounds can increase innervation following denervation. The increase in fiber density does not necessarily indicate regeneration of damaged fibers but may actually arise from collateral and/or terminal sprouting of intact axons. The alternative possibility that the increase in immunoreactivity merely reflects an increase in axonal transport of tyrosine hydroxylase cannot be ruled out. In addition, the possibility of inappropriate innervation of adjacent regions needs to be examined in detail. Whereas functional studies are needed to support the morphological data, a similar study in 6-hydroxydopamine-lesioned rats showed that the FKBP-12 ligand reduced motor disturbance (i.e., amphetamine-induced rotations). Further studies on the ability of FKBP-12 ligands to increase regeneration of CNS axons are warranted to assess their potential for the treatment of human neurodegenerative diseases.

### FK506 Neuronal Actions: Calcineurin-Dependent and Calcineurin-Independent Mechanisms

Taken together, studies employing cyclosporin A and FK506-like compounds demonstrate that calcineurin inhibition is not necessary for FKBP-12 ligands to speed nerve regeneration and implicate a FKBP-pathway distinct from immunophilin-mediated inhibition of calcineurin in the ability of FK506 to increase axonal elongation. Thus, these findings do not support the hypothesis (Gold et al., 1995) that FK506 speeds axonal regeneration by increasing the phosphorylation state of GAP-43 secondary to calcineurin inhibition. However, it is possible that this pathway, by leading to a loss in the dynamics of GAP-43 phosphorylation (Meiri et al., 1991), could

negatively alter nerve regeneration, an explanation for the decreased drug efficacy observed at high (10 mg/kg) daily dosages (*see* FK506 Increases the Rate of Axonal Regeneration in the Rat Sciatic Nerve). Other calcineurin targets can also be discounted as playing a role in FK506's nerve regenerative effect. For example, inhibition of the calcineurin target NOS would lead to a reduction in NO formation (Dawson et al., 1993). Conflicting results have been reported on NO regulation of neurite outgrowth. A preliminary report using sensory neurons indicates that NOS inhibitors increase and, conversely, increased NO formation decreases neurite outgrowth in vitro (Wayne and Skene, 1995). In contrast, a recent paper (Hindley et al., 1997) shows that NO donors increase neurite outgrowth from PC12 cells and primary hippocampal neurons, a result contrary to that expected if NOS inhibition were to play a role in FK506's ability to increase nerve regeneration. Regardless of whether NO positively or negatively impacts neurite outgrowth, a role for the calcineurin target NOS in nerve regeneration is ruled out by studies using FKBP-12 ligands lacking calcineurin inhibition (Steiner et al., 1997a,b; Gold et al., 1997). However, it should be noted that calcineurin inhibition may play a role in mediating other important neuronal properties of FK506. For example, FK506's neuroprotective action against glutamate toxicity in vitro may utilize this pathway (Dawson et al., 1993; Snyder and Sabatini, 1995). In addition, the demonstration that systemic injections of FK506 (Kitamura et al., 1994) and cyclosporin A (Matsuura et al., 1996) protect striatal neurons against depletion of dopamine by MPTP or 6-hydroxydopamine, respectively, strongly suggests that calcineurin inhibition may also mediate this neuroprotective action. Together with the finding that a FKBP-12 ligand increases regeneration of dopaminergic neurons in MPTP-treated mice, these exciting findings suggest that FK506 (or related compounds) may be beneficial in the treatment of Parkinson's disease.

## Putative Mechanisms for FK506's Ability to Increase Nerve Regeneration

The mechanism by which FK506 increases the rate of peripheral nerve regeneration is unknown. Whereas our studies (Wang et al., *in press*) (*see* FKBP-12 Ligand Not Inhibiting Calcineurin Increases Nerve Regeneration) appear to rule out a calcineurin-dependent mechanism, it is possible that FKBP-12 may still be involved, albeit via a different pathway. For example, the FKBP-12 has also been found to demonstrate stable interactions with the two  $\text{Ca}^{2+}$  channels release  $\text{Ca}^{2+}$  from internal stores: the ryanodine (Timerman et al., 1993; Brillantes et al., 1994; Giannini et al., 1995) and the inositol 1,4,5-triphosphate ( $\text{IP}_3$ ) receptors (Cameron et al., 1995a,b); FKBP and calcineurin interact under physiological conditions to modulate  $\text{Ca}^{2+}$  flux in these channels and FK506, by inhibiting calcineurin and preventing receptor dephosphorylation, increases  $\text{Ca}^{2+}$  transport through these channels. The possibility that FK506's ability to stabilize these channels and alter  $\text{Ca}^{2+}$  release (Brillantes et al., 1994; Cameron et al., 1995b) is involved in the drug's regenerative effects (*see* Snyder and Sabatini, 1995) appears to be supported by a very recent preliminary report (Takei et al., 1996) showing that inactivation of the type 1  $\text{IP}_3$  receptor in chick DRG growth cones inhibits neuritic growth. However, the somewhat higher concentrations of FK506 (10–100 nM) necessary to disrupt association of FKBP-12 from the  $\text{IP}_3$  receptor (Cameron et al., 1995a), compared to concentrations that enhance neurite outgrowth in vitro (Lyons et al., 1994; B. G. Gold and M. Zeleny-Pooley, unpublished observation), makes this possibility less attractive.

An alternative mechanism by which FK506 could alter nerve regeneration via FKBP-12 is suggested by the finding that FKBP-12 is a natural ligand for the type 1 receptor for transforming growth factor- $\beta$  (TGF- $\beta$ 1), where it functions as an inhibitor of TGF- $\beta$ 1 receptors (Wang et al., 1994, 1996). Since TGF- $\beta$ 1 stimulates nerve

growth factor (NGF) synthesis by glial cells (Lindholm et al., 1990), and NGF has been suggested to play a role in axonal elongation (Taniuchi et al., 1988), FK506 could increase regeneration indirectly via an increase in NGF. Such a mechanism may be supported by the findings that FK506 appears to increase the sensitivity of PC-12 cells to NGF (Lyons et al., 1994) and by the demonstration that TGF- $\beta$ 1, at similarly low concentrations (i.e., 1 ng/mL), promotes regrowth of injured neurites in vitro (Abe et al., 1996). However, this hypothesis is unattractive given that a role for NGF in axonal regeneration is not supported by in vivo studies (Diamond et al., 1992); in fact, we have shown recently that delivery of NGF to the neuronal cell body (by intrathecal infusion) delays regeneration (Gold, 1997). Furthermore, FKBP-12 has been shown to be dispensable for TGF- $\beta$  signaling (Chamg et al., 1996). In addition, as in the case of the IP<sub>3</sub> receptor (*see* preceding paragraph), much higher concentrations of FK506 ( $\mu$ M) are needed to disrupt association of FKBP-12 from the TGF- $\beta$ 1 receptor (Wang et al., 1994) relative to those that stimulate neurite outgrowth in vitro (Lyons et al., 1994); B. G. Gold and M. Zeleny-Pooley, unpublished observation.

An unexplored area is the possible involvement of FKBP-12 in mediating FK506's nerve regenerative effect. Most interesting is the FK506-FKBP-52 complex, which does not inhibit calcineurin (*see* Snyder and Sabatini, 1995). Instead, human FKBP-52 (rabbit FKBP-59) (human FKBP-52) (Tai et al., 1992) has been identified as a heat-shock protein (hsp-56) (Sanchez, 1990) and, together with the hsp-90 and hsp-70, this novel immunophilin comprises a component of a subclass of glucocorticoid receptor complexes (Owens-Grillo, et al., 1995; Perdew and Whitelaw, 1991; Lebeau et al., 1992; Czar et al., 1994); FK506 does not alter glucocorticoid signaling pathways but may produce conformational changes in the complex (Ratajczak and Carrello, 1996). In this context, we have found that subcutaneous injection of FK506 (10 mg/kg/d, for 2 wk) increases expression of hsp-70 (as shown by immunocytochemistry) in selected neurons in

the brain (including the cortex, hippocampus, and amygdala), and in the spinal cord and dorsal root ganglion (DRG) (Goto and Singer, 1994). It may therefore be relevant that the glucocorticoid dexamethasone has been found to increase GAP-43 mRNA levels in regenerating hypoglossal neurons (Yao and Kiyama, 1995). Moreover, it has recently been shown that FK506 potentiates the potency of steroid hormones (Kralli and Yamamoto, 1996). Thus, our (B. G. Gold, J. Y. Yew, and M. Zeleny-Pooley, unpublished observations) recent demonstration that FK506 increases GAP-43 mRNA levels in DRG neurons may support an involvement of this pathway in FK506's ability to increase nerve regeneration.

Consistent with our nerve regenerative studies (*see* Cyclosporin A Does Not Increase Axonal Regeneration Rate), the ability of FK506 to protect hippocampal CA1 and cerebral neurons against transient global ischemia is not shared by cyclosporin A (Yagita et al., 1996). This suggests that a calcineurin-independent mechanism also underlies FK506's ability to provide neuroprotection against ischemic injury. The recent demonstration that global ischemia induces an alteration in the ryanodine receptor (Nozaki et al., 1996) suggests that FK506 via the FKBP-12 may protect against damage via binding to the FKBP-12. However, the necessity of using repeated drug administration (Drake et al., 1996) suggests that simple inhibition of NO may not fully explain the neuroprotective actions of FK506. FK506's ability to protect against focal cerebral ischemia (Sharkey and Butcher, 1994; Ide et al., 1996; Tokime et al., 1996) may therefore involve additional FKBP pathways, perhaps mediated by hsp-70 induction via glucocorticoid receptors. Whether this pathway also plays a role in the ability of FK506 to protect against ischemic damage is an important area for future work.

Why FK506 exhibits a reduction in efficacy for axonal regeneration at high (10 mg/kg) dosages is unknown. Interestingly, a bell-shaped dose-response has been observed for other agents that increase regeneration. For example, the melanocortins ( $\alpha$ -MSH), insulin-like growth factor-I, and ACTH demonstrate a

similar decrease in efficacy beyond a certain optimal dosage (Bijlsma et al., 1981; Contreras et al., 1995; Bijlsma et al., 1983); however, for the latter two agents, hypoglycemia and corticosteroid activity, respectively, may be responsible for the reduced regenerative efficacy of high dosages (Bijlsma et al., 1981; Contreras et al., 1995). For the melanocortins, it has been suggested (Joosten et al., 1996) that the bell-shaped dose-response is caused by the action of multiple melanocortin receptors (Hol et al., 1995). Whereas alternative mechanisms are indeed possible, the reduced efficacy of higher (10 mg/kg) dosages of FK506 suggests that multiple FKBP (Snyder and Sabatini, 1995; Sigal and Dumont, 1992; Cardenas et al., 1995; Wiederrecht et al., 1993) or FKBP-mediated pathways (acting in opposition) underlie the ability of FK506 to alter nerve regeneration. For example, whereas the results presented in this review indicate that FK506 speeds regeneration via a calcineurin-independent pathway, calcineurin inhibition (which may be less sensitive) would be expected to impair regeneration (Chang et al., 1995) leading to a reduction in efficacy at higher (10 mg/kg) dosages of FK506.

## Summary and Future Directions

Taken together, it appears that distinct mechanisms underlie the immunosuppressant (calcineurin-dependent) and nerve regenerative (calcineurin-independent) properties of FK506. Based upon structural analysis of FK506-FKBP interactions (Griffith et al., 1995; Itoh et al., 1995a,b), it has been possible to separate these properties and design new FKBP ligands (Armistead et al., 1995; Batchelor et al., 1994; Shuker et al., 1996) that do not inhibit calcineurin. Of equal importance for drug development is the possibility that different pathways underlie FK506's nerve regenerative (Wang et al., 1997; Steiner et al., 1997a,b; Gold et al., 1997) and neurotoxic properties (Lopez et al., 1991; Mueller et al., 1994; Wijdicks et al., 1995; Bronster et al., 1995; Vincenti et al., 1996). The latter have been linked mechanistically to the

immunosuppressant properties of FK506 via calcineurin inhibition (Dumont et al., 1992); calcineurin inhibition also has been shown to underlie development of some pathological hallmarks of Alzheimer's disease (Ladner et al., 1996). Thus, it is possible that FK506's neurotoxic properties will not be shared by these new (noncalcineurin-binding) FKBP ligands.

It should be apparent, however, that our knowledge is critically lacking in regard to the underlying molecular mechanism(s) by which FKBP ligands produce their neuronal effects. Of paramount importance will be identification of the molecular targets for FKBP ligand-FKBP complexes. This information should become available in the near future given the recent development of innovative new techniques such as the three-hybrid system (Licitra and Liu, 1996) to "fish" for such targets. It will then be necessary to sort out which of these targets are physiologically relevant for promoting nerve regeneration. These studies should also shed new light on the neuron's intrinsic mechanism(s) regulating axonal elongation. Regardless of the underlying mechanism involved, the development of novel FKBP ligands (Navia and Chaturvedi, 1996) may lead to the generation of new drugs for the treatment of human nerve injuries.

## Acknowledgments

I thank Michelle Zeleny-Pooley and Dan Austin for preparation of figures, Fujisawa Pharmaceuticals, Osaka, Japan for the generous gift of FK506, and Vertex Pharmaceuticals, Cambridge, MA for providing compound V-10,367. I especially would like to thank David Armistead (Vertex Pharmaceuticals) for sharing information on the chemistry of V-10,367 and for helpful discussions concerning the manuscript, and Solomon Snyder (Johns Hopkins University) for providing advanced copies of two manuscripts. I am indebted to Peter S. Spencer for his insightful and critical review of the manuscript. This paper is dedicated to my brother, Michael S. Gold (University of New Mexico).

## References

- Abe K., Chu P.-J., Ishihara A., and Saito H. (1996) Transforming growth factor- $\beta$ 1 promotes re-elongation of injured axons of cultured rat hippocampal neurons. *Brain Res.* **723**, 206–209.
- Alnemri E. S., Fernandes-Alnemri T., Nelki D. S., Dudley K., DuBois G. C., and Litwack G. (1993) Overexpression, characterization, and purification of a recombinant mouse immunophilin FKBP-52 and identification of an associated phosphoprotein. *Proc. Natl. Acad. Sci. USA* **90**, 6839–6843.
- Angelov D. N., Neiss V. F., Streppel M., Andermahr J., Mader K., and Stennert E. (1996) Nimodipine accelerates axonal sprouting after surgical repair of rat facial nerve. *J. Neurosci.* **16**, 1041–1048.
- Armistead D. M., Badia M. C., Deininger D. D., Duffy J. P., Saunders J. O., Tung R. D., Thomson J. A., DeCenzo M. T., Futer O., Livingston D. J., Murcko M. A., Yamashita M. M., and Navia M. A. (1995) Design, synthesis and structure of non-macrocyclic inhibitors of FKBP12, the major binding protein for the immunosuppressant FK506. *Acta Cryst.* **51**, 522–528.
- Bächinger H. P., Morris N. P., and Davis J. M. (1993) Thermal stability and folding of the collagen triple helix and the effects of mutations in osteogenesis imperfecta on the triple helix of type I collagen. *Am. J. Med. Genet.* **45**, 152–162.
- Banker G. A. and Cowan W. M. (1977) Rat hippocampal neurons in dispersed cell culture. *Brain Res.* **126**, 397–425.
- Batchelor M. J., Gillespie R. J., Golec J. M. C., Hedgecock C. J. R., Jones S. D., and Murdoch R. (1994) Total synthesis of close analogues of the immunosuppressant FK506. *Tetrahedron* **50**, 809–825.
- Baughman G., Wiederrecht G. J., Campbell N. F., Martin M. M., and Bourgeois S. (1995) FKBP51, a novel T-cell-specific immunophilin capable of calcineurin inhibition. *Mol. Cell Biol.* **15**, 4395–4402.
- Bennett P. C., Zhao W., Lawen A., and Ng K. T. (1996) Cyclosporin A, an inhibitor of calcineurin, impairs memory formation in day-old chicks. *Brain Res.* **730**, 107–117.
- Beuche W. and Friede R. L. (1984) The role of non-resident cells in Wallerian degeneration. *J. Neurocytol.* **13**, 767–796.
- Bierer B. E., Somers P. K., Wandless T. J., Burakoff S. J., and Schreiber S. L. (1990) Probing immunosuppressant action with a nonnatural immunophilin ligand. *Science* **250**, 556–559.
- Bijlsma W. A., Jennekens W. A., Schotman P., and Gispen W. H. (1984) Neurotrophic factors and regeneration in the peripheral nervous system. *Psychoneuroendocrinology* **9**, 199–215.
- Bijlsma W. A., Jennekens F. G. I., Schotman P., and Gispen W. H. (1983) Stimulation of ACTH-10 of nerve fiber regeneration following sciatic nerve crush. *Muscle Nerve* **6**, 104–112.
- Bijlsma W. A., Jennekens F. G. I., Schotman P., and Gispen W. H. (1981) Effects of corticotrophin (ACTH) on recovery of sensorimotor function in the rat: structure-activity study. *Eur. J. Pharmacol.* **76**, 73–79.
- Bixby J. L. and Jhabvala P. (1993) Tyrosine phosphorylation in early embryonic growth cones. *J. Neurosci.* **13**, 3421–3432.
- Brück W. and Friede R. L. (1990) Anti-macrophage CR3 antibody blocks myelin phagocytosis by macrophages *in vitro*. *Acta Neuropath.* **103**, 182–187.
- Brillantes A. M. B., Ondrias K., Scott A., Kobrinsky E., Ondriasova E., Moschella M. C., Jayaraman T., Landers M., Ehrlich B. E., and Marks A. R. (1994) Stabilization of calcium release channel (ryanodine receptor) function by FK506-binding protein. *Cell* **77**, 513–523.
- Bronster D. J., Yonover P., Stein J., Scelsa S. N., Miller C. M., and Sheiner P. A. (1995) Demyelinating sensorimotor polyneuropathy after administration of FK506. *Transplantation* **59**, 1066–1068.
- Butcher S. P., Henshall D. C., Teramura Y., Iwasaki K., and Sharkey J. (1997) Neuroprotective actions of FK506 in experimental stroke: *in vivo* evidence against an anti-excitotoxic mechanism. *J. Neurosci.* **17**, 6939–6946.
- Cameron A. M., Steiner J. P., Sabatini D. M., Kaplin A. I., Walensky L. D., and Snyder S. H. (1995a) Immunophilin FK506 binding protein associated with inositol 1,4,5-trisphosphate receptor modulates calcium flux. *Proc. Natl. Acad. Sci. USA* **92**, 1784–1788.
- Cameron A. M., Steiner J. P., Roskams A. J., Ali S. M., Ronnett G. V., and Snyder S. H. (1995b) Calcineurin associated with the inositol 1,4,5-trisphosphate receptor-FKBP12 complex modulates  $Ca^{2+}$  flux. *Cell* **83**, 463–472.
- Cardenas M. E., Zhu D., and Heitman J. (1995) Molecular mechanisms of immunosuppression by cyclosporine, FK506, and rapamycin. *Curr. Opin. Nephrol. Hypertens.* **4**, 472–477.
- Chang H. Y., Takei K., Sydor A. M., Born T., Rusnak F., and Jay D. G. (1995) Asymmetric retraction of

- growth cone filopodia following focal inactivation of calcineurin. *Nature* **376**, 686–690.
- Charng M.-J., Kinnunen P., Hawker J., Brand T., and Schneider M. D. (1996) FKBP-12 recognition is dispensable for signal generation by type I transforming growth factor- $\beta$  receptors. *J. Biol. Chem.* **271**, 22941–22944.
- Chen S. and Bisby M. A. (1993) Impaired motor axon regeneration in the C57BL/Ola mouse. *J. Comp. Neurol.* **333**, 449–454.
- Compton L. A., Davis J. M., MacDonald J. R., and Bächinger H. P. (1992) Structural and function characterization of *Escherichia coli* peptidyl-prolyl *cis-trans* isomerases. *Eur. J. Biochem.* **206**, 927–934.
- Contreras P. C., Steffler C., Yu E., Callison K., Stong D., and Vaught J. L. (1995) Systemic administration of rhIGF-I enhanced regeneration after sciatic nerve crush in mice. *J. Pharmacol. Exp. Ther.* **274**, 1443–1449.
- Cosenza C. A., Cramer D. V., and Makowka L. (1994) The use of immunosuppressive agents in animal models of experimental transplantation, in *Handbook of Animal Models in Transplantation Research* (Cramer D. V., Posesta L., and Makowka L. eds.), CRC Press, Boca Raton, FL, pp. 331–337.
- Czar M. J., Owens-Grillo J. K., Yem A. W., Leach K. L., Deibel M. R., Jr., Welsh M. J., and Pratt W. B. (1994) The hsp56 immunophilin component of untransformed steroid receptor complexes is localized both to microtubules in the cytoplasm and to the same nonrandom regions within the nucleus as the steroid receptor. *Mol. Endocrin.* **8**, 1731–1741.
- Dawson T. M., Steiner J. P., Dawson V. L., Dinerman J. L., Uhl G. R., and Snyder S. H. (1993) Immunosuppressant FK506 enhances phosphorylation of nitric oxide synthase and protects against glutamate neurotoxicity. *Proc. Natl. Acad. Sci. USA* **90**, 9808–9812.
- De Koning P. and Gispén W. H. (1987a) A rationale for the use of melanocortins in neural damage, in *Pharmacological Approaches to the Treatment of Brain and Spinal Cord Injury* (Stein D. G. and Sabel C., eds.), Plenum, New York, pp. 233–258.
- De Koning P. and Gispén W. H. (1987b) Org. 2766 improves functional and electrophysiological aspects of regenerating sciatic nerve in the rat. *Peptides* **8**, 415–422.
- Diamond J., Foerster A., Holmes M., and Coughlin M. (1992) Sensory nerves in adult rats regenerate and restore sensory function to the skin independently of endogenous NGF. *J. Neurosci.* **12**, 1467–1476.
- Drake M., Friberg H., Boris-Möller F., Sakata K., and Wieloch T. (1996) The immunosuppressant FK506 ameliorates ischaemic damage in the rat brain. *Acta Physiol. Scand.* **158**, 155–159.
- Dumont F. J., Staruch M. J., Koprak S. L., Siekierka J. J., Lin C. S., Harrison R., Sewell T., Kindt V. M., Beattie T. R., Wyvratt M., and Sigal N. H. (1992) The immunosuppressive and toxic effects of FK-506 are mechanistically related: pharmacology of a novel antagonist of FK-506 and rapamycin. *J. Exp. Med.* **176**, 751–760.
- Edwards P. M., Verhaagen J., Spierings T., Schotman P., Jennekens F. G. I., and Gispén W. H. (1985) The effect of ACTH<sub>4-10</sub> on protein synthesis, actin and tubulin during regeneration. *Brain Res. Bull.* **15**, 267–272.
- Ekström P. A. R. and Tomlinson D. R. (1990) Impaired nerve regeneration in streptozotocin-diabetic rats is improved by treatment with gangliosides. *Exp. Neurol.* **109**, 200–203.
- Ekström P. A. R., Kanje M., and Skottner A. (1989) Nerve regeneration and serum levels of insulin-like growth factor-I in rats with streptozotocin-induced insulin deficiency. *Brain Res.* **496**, 141–147.
- Figueroas A., Morales-Olivas F. J., and Capella D. (1992) Bovine gangliosides and acute motor polyneuropathy. *Brit. Med. J.* **305**, 1330–1331.
- Freeman D. A., Klee C. B., Bierer B. E., and Burakoff S. J. (1992) Calcineurin phosphatase activity in T lymphocytes is inhibited by FK506 and cyclosporin A. *Proc. Natl. Acad. Sci. USA* **89**, 3686–3690.
- Friede R. L. and Brück W. (1993) Macrophage functional properties during myelin degradation. *Adv. Neurol.* **59**, 327–336.
- Funauchi M., Haruta H., and Tsumoto T. (1994) Effects of an inhibitor for calcium/calmodulin-dependent protein phosphatase, calcineurin, on induction of long-term potentiation in rat visual cortex. *Neurosci. Res.* **19**, 269–278.
- Fung J. J., Eliasziw M., Todo S., Jain A., Demetris A. J., McMichael J. P., Starzl T. E., Meier P., and Donner A. (1996) The Pittsburgh randomized trial of tacrolimus compared to cyclosporine for hepatic transplantation. *J. Am. Coll. Surg.* **183**, 117–125.
- Gallai V., Mazzotta G., Montesi S., Sarchielli P., and Del Gatto F. (1992) Effects of uridine in the treatment of diabetic neuropathy: an electrophysiological study. *Acta Neurol. Scand.* **86**, 3–7.



- Giannini G., Conti A., Mammarella S., Scrobogna M., and Sorrentino V. (1995) The ryanodine receptor/calcium channel genes are widely and differentially expressed in murine brain and peripheral tissues. *J. Cell Biol.* **128**, 893–904.
- Gispen W. H., Sporel-Özkat E., Duckers E., and Edwards P. M. (1991) The  $\text{Ca}^{2+}$  antagonist nifedipine and recovery of function following peripheral nerve damage in the rat. *Neurosci. Res. Commun.* **8**, 175–184.
- Gold B. G. (1997) Axonal regeneration of sensory neurons is delayed by continuous intrathecal infusion of nerve growth factor. *Neuroscience* **76**, 1153–1158.
- Gold B. G., Zeleny-Pooley M., Wang M.-S., Chaturvedi P., and Armistead D. M. (1997) A non-immunosuppressant FKBP-12 ligand increases nerve regeneration. *Exp. Neurol.* **147**, 269–278.
- Gold B. G., Katoh K., and Storm-Dickerson T. (1995) The immunosuppressant FK506 increases the rate of axonal regeneration in rat sciatic nerve. *J. Neurosci.* **15**, 7505–7516.
- Gold B. G., Storm-Dickerson T., and Austin D. R. (1994) The immunosuppressant FK506 increases functional recovery and nerve regeneration following peripheral nerve injury. *Restor. Neurol. Neurosci.* **6**, 287–296.
- Gold B. G., Storm-Dickerson T., Austin D. R., and Katoh K. (1993) FK506, an immunosuppressant, increases functional recovery and axonal regeneration in the rat following axotomy of the sciatic nerve. *Soc. Neurosci. Abstr.* **19**, 1316.
- Goldberger M. M., Murray M., and Tessler A. (1993) Sprouting and regeneration in the spinal cord: Their roles in recovery of function after spinal injury, in *Neuroregeneration* (Gorio A., ed.), Raven, New York, pp. 241–264.
- Gorio A., DiGiulio A. M., and Mantegazza P. (1993) Pharmacology of neuronal regeneration, in *Neuroregeneration* (Gorio A., ed.), Raven, New York, pp. 289–320.
- Goto S. and Singer W. (1994) Laminar and columnar organization of immunoreactivity for calcineurin, a calcium- and calmodulin-regulated protein phosphatase, in monkey striate cortex. *Cereb. Cortex* **4**, 636–645.
- Griffin J. W., George R., and Ho T. (1993) Macrophage systems in peripheral nerves: a review. *J. Neuropathol. Exp. Neurol.* **52**, 553–560.
- Griffith J. P., Kim J. L., Kim E. E., Sintchank M. D., Thomson J. A., Fitzgibbon M. J., Fleming M. A., Caron P. R., Hsiao K., and Navia M. A. (1995) X-ray structure of calcineurin inhibited by the immunophilin-immunosuppressant FKBP12-FK506 complex. *Cell* **82**, 507–522.
- Harding M. W., Galat A., Uehling D. E., and Schreiber S. L. (1989) A receptor for the immunosuppressant FK506 is a *cis-trans* peptidyl-prolyl isomerase. *Nature* **341**, 758–760.
- Hindley S., Juurlink B. H. J., Gusbers J. W., Middlemiss P. J., Herman M. A. R., and Rothbone M. P. (1997) Nitric oxide donors enhance neurotrophin-induced neurite outgrowth through a cGMP-dependent mechanism. *J. Neurosci. Res.* **47**, 427–439.
- Hodgkiss J. P. and Kelly J. S. (1995) Only “de novo” long-term depression (LTD) in the rat hippocampus in vitro is blocked by the same low concentration of FK506 that blocks LTD in the visual cortex. *Brain Res.* **705**, 241–246.
- Hoffman A. L., Makowka L., Banner B., Cai X., Cramer D. V., Pascualone A., Todo S., and Starzl T. E. (1990) The use of FK-506 for small intestine allotransplantation: inhibition of acute rejection and prevention of fatal graft-versus-host disease. *Transplantation* **49**, 483–490.
- Hol E., Gispen W. H., and Bär P. R. (1995) ACTH-related peptides: Receptors and signal transduction systems involved in their neurotrophic and neuroprotective actions. *Peptides* **16**, 979–993.
- Ide T., Morikawa E., and Kirino T. (1996) An immunosuppressant, FK506, protects hippocampal neurons from forebrain ischemia in the Mongolian gerbil. *Neurosci. Lett.* **204**, 157–160.
- Ikegami S., Kato A., Kudo Y., Kuno T., Ozawa F., and Inokuchi K. (1996) A facilitatory effect on the induction of long-term potentiation in vivo by chronic administration of antisense oligodeoxynucleotides against catalytic subunits of calcineurin. *Mol. Brain Res.* **41**, 183–191.
- Itoh S., DeCenzo M. T., Livingston D. J., Pearlman D. A., and Navia M. A. (1995a) Conformation of FK506 in x-ray structures of its complexes with human recombinant FKBP12 mutants. *Bioorganic Medicinal Chem. Lett.* **5**, 1983–1988.
- Itoh S. and Navia M. A. (1995b) Structure comparison of native and mutant human recombinant FKBP12 complexes with the immunosuppressant drug FK506 (tacrolimus). *Protein Sci.* **4**, 2261–2268.
- Jones K. J. (1993) Gonadal steroids as promoting factors in axonal regeneration. *Brain Res. Bull.* **30**, 491–498.



- Joosten E. A. J., Verhaagh S., Martin D., Robe P., Franzen R., Hooiveld M., Doornbos R., Bär P. R., and Moonen G. (1996) Alpha-MSH stimulates neurite outgrowth of neonatal rat corticospinal neurons *in vitro*. *Brain Res.* **736**, 91–98.
- Känge M., Skottner A., Sjöberg J., and Lundborg G. (1989) Insulin-like growth factor I (IGF-I) stimulates regeneration of the rat sciatic nerve. *Brain Res.* **486**, 396–398.
- Kantor D. B., Lanzrein M., Stary S. J., Sandoval G. M., Smith W. B., Sullivan B. M., Davidson N., and Schuman E. M. (1996) A role for endothelial NO synthase in LTP revealed by adenovirus-mediated inhibition and rescue. *Science* **274**, 1744–1748.
- Kino T. and Goto T. (1993) Discovery of FK-506 and update. *Ann. NY Acad. Sci.* **685**, 13–21.
- Kino T., Hatanaka H., Hashimoto M., and Nishiyama H. (1987a) FK-506, a novel immunosuppressant isolated from a streptomyces. I. Fermentation, isolation, and physiochemical and biological characteristics. *J. Antibiot.* **40**, 1249–1255.
- Kino T., Hatanaka H., Miyata S., Inamura N., Nishiyama M., Yajima T., Goto T., Okuhara M., Kohsaka M., Aoki H., and Ochiai T. (1987b) FK-506, a novel immunosuppressant isolated from a streptomyces. II. Immunosuppressive effect of FK-506 *in vitro*. *J. Antibiot.* **40**, 1256–1265.
- Kitamura Y., Itano Y., Kubo T., and Nomura Y. (1994) Suppressive effect of FK-506, a novel immunosuppressant, against MPTP-induced dopamine depletion in the striatum of young C57BL/6 mice. *J. Neuroimmunol.* **50**, 221–224.
- Klee C. B. (1991) Concerted regulation of protein phosphorylation and dephosphorylation by calmodulin. *Neurochem. Res.* **16**, 1059–1065.
- Klintmalm G. (1994) The U.S. Multicenter FK 506 Liver Study Group: a comparison of tacrolimus (FK 506) and cyclosporine for immunosuppression in liver transplantation. *N. Engl. J. Med.* **331**, 1110–1115.
- Kralli A. and Yamamoto K. R. (1996) An FK506-sensitive transporter selectively decreases intracellular levels and potency of steroid hormones. *J. Biol. Chem.* **271**, 17152–17156.
- Kuromi H., Yoshihara M., and Kidokoro Y. (1997) An inhibitory role of calcineurin in endocytosis of synaptic vesicles at nerve terminals of *Drosophila* larvae. *Neurosci. Res.* **27**, 101–113.
- Ladner C. J., Czech J., Maurice J., Lorens S. A., and Lee J. M. (1996) Reduction of calcineurin enzymatic activity in Alzheimer's disease: correlation with neuropathologic changes. *J. Neuropathol. Exp. Neurol.* **55**, 924–931.
- Landi G., D'Alessandro R. D., Dossi B. C., Ricci S., Simone I. L., and Ciccone A. (1993) Guillain-Barré syndrome after exogenous gangliosides in Italy. *Brit. Med. J.* **307**, 1463–1464.
- Lebeau M. C., Massol N., Herrick J., Faber L. E., Renoin M. J., Radanyi C., and Baulieu E. E. (1992) P59, an hsp 90-binding protein. Cloning and sequencing of its cDNA and preparation of a peptide-directed polyclonal antibody. *J. Biol. Chem.* **267**, 4281–4284.
- Letteron P., Fouin-Fortunet H., Tinel M., Danan G., Belghiti J., and Pessayre D. (1984) Mechanism for isaxoxine hepatitis. I. Metabolic activation by mouse and human cytochrome P-450. *J. Pharmacol. Exp. Ther.* **229**, 845–850.
- Lewin G. R. (1995) Neurotrophic factors and pain. *Neuroscience* **7**, 227–232.
- Licitra E. J. and Liu J. O. (1996) A three-hybrid system for detecting small ligand-protein receptor interactions. *Proc. Natl. Acad. Sci. USA* **93**, 12817–12821.
- Lindholm D., Hengerer B., Zafra F., and Thoenen H. (1990) Transforming growth factor- $\beta$ 1 stimulates expression of nerve growth factor in the rat CNS. *NeuroReport* **1**, 9–12.
- Liu J., Farmer J. D., Lane W. S., Friedman J., Weissman I., and Schreiber S. L. (1991) Calcineurin is a common target of cyclophilin-cyclosporin A and FKBP-FK506 complexes. *Cell* **66**, 807–815.
- Liu Y. and Storm D. R. (1990) Regulation of free calmodulin levels by neuromodulin: neuron growth and regeneration. *Trends Pharmacol. Sci.* **11**, 107–111.
- Lopez O. L., Martinez A. J., and Torre-Cisneros J. (1991) Neuropathologic findings in liver transplantation: a comparative study of cyclosporine and FK506. *Transplant. Proc.* **23**, 3181–3182.
- Ludwin S. K. and Bisby M. A. (1992) Delayed Wallerian degeneration in the central nervous system of Ola mice: an ultrastructural study. *J. Neurol. Sci.* **109**, 140–147.
- Lyons W. E., Steiner J. P., Snyder S. H., and Dawson T. M. (1995) Neuronal regeneration enhances the expression of the immunophilin FKBP-12. *J. Neurosci.* **15**, 2985–2994.
- Lyons W. E., George E. B., Dawson T. M., Steiner J. P., and Snyder S. H. (1994) Immunosuppressant FK506 promotes neurite outgrowth in cultures of PC12 cells and sensory ganglia. *Proc. Natl. Acad. Sci. USA* **91**, 3191–3195.

- Lyons W. E., Steiner J. P., Dawson T. M., and Snyder S. H. (1992) Increased expression of FK-506 binding protein during peripheral nerve regeneration. *Soc. Neurosci. Abstr.* **18**, 604.
- Malen P. L. and Chapman P. F. (1997) Nitric oxide facilitates long-term potentiation, but not long-term depression. *J. Neurosci.* **17**, 2645–2651.
- Matsuura K., Kabuto H., Makino H., and Ogawa N. (1996) Cyclosporin A attenuates degeneration of dopaminergic neurons induced by 6-hydroxydopamine in the mouse brain. *J. Neuroimmunol.* **733**, 101–104.
- McDiarmid S. V., Busuttil R. W., Ascher N. L., Burdick A. J., D'Alessandro A. M., Esquivel C., Kalayoglu M., Klein A. S., Marsh J. W., Miller C. M., Schwartz M. E., Shaw B. W., and So S. K. (1995) FK506 (tacrolimus) compared with cyclosporine for primary immunosuppression after pediatric liver transplantation. *Transplantation* **59**, 530–536.
- Meiri K. F., Bickerstaff L. E., and Schwob J. E. (1991) Monoclonal antibodies show that kinase C phosphorylation of GAP-43 during axonogenesis is both spatially and temporally restricted in vivo. *J. Cell Biol.* **112**, 991–1005.
- Moriwaki A., Lu Y.-F., Hayashi Y., Tomizawa K., Tokuda M., Itano T., Hatase O., and Matsui H. (1996) Immunosuppressant FK506 prevents mossy fiber sprouting induced by kindling stimulation. *Neurosci. Res.* **25**, 191–194.
- Mueller A. R., Platz K.-P., Bechstein W.-O., Schattenfroh N., Stoltenburg-Didinger G., Blumhardt G., Christe W., and Neuhaus P. (1994) Neurotoxicity after orthotopic liver transplantation. A comparison between cyclosporine and FK506. *Transplantation* **58**, 155–169.
- Mukai H., Kuno T., Chang C.-D., Lane B., Luly J. R., and Tanaka C. (1993) FKBP12-FK506 complex inhibits phosphatase activity of two mammalian isoforms of calcineurin irrespective of their substrates or activation mechanisms. *J. Biochem.* **113**, 292–298.
- Murad F. (1990) Drugs used for the treatment of angina: organic nitrates, calcium-channel blockers, and  $\beta$ -adrenergic antagonists, in *Goodman and Gilman's: The Pharmacological Basis of Therapeutics* (Gilman A. G., Rall T., Nies A. S., and Taylor P. eds.), Pergamon, New York, pp. 764–783.
- Murase N., Todo S., Lee P.-H., Chapman F., Nalesnik M. A., Makowka L., and Starzl T. E. (1987) Heterotopic heart transplantation in the rat receiving FK-506 alone or with cyclosporine. *Transplant. Proc.* **19**, 71–75.
- Navia M. A. and Chaturvedi P. R. (1996) Design principles for orally bioavailable drugs. *DDT* **1**, 179–189.
- Neuhaus P., Pichlmayr R., and Williams R. (1994) The European FK 506 Multicentre Liver Study Group: randomised trial comparing tacrolimus (FK 506) and cyclosporin in prevention of liver allograft rejection. *Lancet* **344**, 423–428.
- Nozaki H., Tanaka K., Gomi S., Mihara B., Nogawa S., Nagata E., Kondo T., and Fukuuchi Y. (1996) Alteration of ryanodine receptor in the hippocampus CA1 after hemispheric cerebral ischemia. *Neurochem. Res.* **21**, 975–982.
- O'Dell T. J., Hawkins R. D., Kandel E. R., and Arancio O. (1991) Tests of the roles of two diffusible substances in long-term potentiation: evidence for nitric oxide as a possible early retrograde messenger. *Proc. Natl. Acad. Sci. USA* **88**, 11285–11289.
- Ochiai T., Sakamoto K., Nagata M., Nakajima K., Goto T., Hori S., Kenmochi T., Nakagori T., Asano T., and Isono K. (1988) Studies on FK-506 in experimental organ transplantation. *Transplant. Proc.* **20**, 209–214.
- Ochiai T., Nakajima K., Nagata M., Suzuki T., Asano T., Uematsu T., Goto T., Hori S., Kenmochi T., Nakagori T., and Isono K. (1987) Effect of a new immunosuppressive agent, FK 506, on heterotopic cardiac allotransplantation in the rat. *Transplant. Proc.* **19**, 1284–1286.
- Odom A., Del Poeta M., Perfect J., and Heitman J. (1997) The immunosuppressant FK506 and its nonimmunosuppressive analog L-685,818 are toxic to *Cryptococcus neoformans* by inhibition of a common target protein. *Antimicrob. Agents Chemother.* **41**, 156–161.
- Owens-Grillo J. K., Hoffmann K., Hutchison K. A., Yem A. W., Deibel M. R., Jr., Handschumacher R. E., and Pratt W. B. (1995) The cyclosporin A-binding immunophilin CyP-40 and the FK506-binding immunophilin hsp56 bind to a common site on hsp90 and exist in independent cytosolic heterocomplexes with the untransformed glucocorticoid receptor. *J. Biol. Chem.* **270**, 20479–20488.
- Perdew G. H. and Whitelaw M. L. (1991) Evidence that the 90-kDa heat shock protein (HSP90) exists in cytosol in heteromeric complexes containing HSP70 and three other proteins with  $M_r$  of 63,000, 56,000, and 50,000. *J. Biol. Chem.* **266**, 6708–6713.

- Perry V. H., Brown M. C., Lunn E. R., Tree P., and Gordon S. (1990a) Evidence that very slow Wallerian degeneration in C57BL/Ola mice is an intrinsic property of the peripheral nerve. *Eur. J. Neurosci.* **2**, 802–808.
- Perry V. H., Lunn E. R., Brown M. C., Cahusac S., and Gordon S. (1990b) Evidence that the rate of Wallerian degeneration is controlled by a single autosomal dominant gene. *Eur. J. Neurosci.* **2**, 408–413.
- Ratajczak T. and Carrello A. (1996) Cyclophilin 40 (CyP-40), mapping of its hsp90 binding domain and evidence that FKBP52 competes with CyP-40 for hsp90 binding. *J. Biol. Chem.* **271**, 2961–2965.
- Roth G. A., Spada V., Hamill K., and Bornstein M. B. (1995) Insulin-like growth factor I increases myelination and inhibits demyelination in cultured organotypic nerve tissue. *Dev. Brain Res.* **88**, 102–108.
- Sakai K., Date I., Yoshimoto Y., Arisawa T., Nakashima H., Furuta T., Asari S., and Ohmoto T. (1991) The effect of a new immunosuppressive agent, FK-506, on xenogeneic neural transplantation in rodents. *Brain Res.* **565**, 167–170.
- Sanchez E. R. (1990) Hsp56: a novel heat shock protein associated with untransformed steroid receptor complexes. *J. Biol. Chem.* **265**, 22067–22070.
- Sánchez E. R. and Ning Y.-M. (1996) Immunophilins, heat shock protein, and glucocorticoid receptor actions in vivo. *Meth. Enzymol.* **9**, 188–200.
- Schonhofer P. S. (1991) Guillain-Barré syndrome and parenteral gangliosides. *Lancet* **338**, 757.
- Schreiber S. L. and Crabtree G. R. (1992) The mechanism of action of cyclosporin A and FK506. *Immunol. Today* **13**, 136–142.
- Schreiber S. L. (1991) Chemistry and biology of the immunophilins and their immunosuppressive ligands. *Science* **251**, 283–287.
- Schuman E. M. and Madison D. V. (1991) A requirement for the intercellular messenger nitric oxide in long-term potentiation. *Science* **254**, 1503–1506.
- Sebille A., Saint-André P., and Hugelin A. (1982) Acceleration by isaxonine of muscle reinnervation: electrophysiological demonstration in leprotic neuropathy. *Nouv. Presse. Med.* **11**, 1278–1282.
- Shapiro R., Fung J. J., Jain A. B., Parks P., Todo S., and Starzl T. E. (1990) The side effects of FK506 in humans. *Transplant. Proc.* **22**, 35–36.
- Sharkey J. and Butcher S. P. (1994) Immunophilins mediate the neuroprotective effects of FK506 in focal cerebral ischaemia. *Nature* **371**, 336–339.
- Shibuki K. and Okada D. (1996) Endogenous nitric oxide release required for long-term synaptic depression in the cerebellum. *Nature* **349**, 326–328.
- Shuker S. B., Hajduk P. J., Meadows R. P., and Fesik S. W. (1996) Discovering high-affinity ligands for proteins: SAR by NMR. *Science* **274**, 1531–1534.
- Siekierka J. J., Wiederrecht G., Greulich H., Boulton D., Hung S. H. Y., Cryan J., Hodges P. J., and Sigal N. H. (1990) The cytosolic-binding protein for the immunosuppressant FK-506 is both a ubiquitous and highly conserved peptidyl-prolyl *cis-trans* isomerase. *J. Biol. Chem.* **265**, 21011–21015.
- Siekierka J. J., Hung S. H. Y., Poe M., Lin C. S., and Sigal N. H. (1989) A cytosolic binding protein for the immunosuppressant FK506 has peptidyl-prolyl isomerase activity but is distinct from cyclophilin. *Nature* **341**, 755–757.
- Sigal N. H. and Dumont F. J. (1992) Cyclosporin A, FK-506, and rapamycin: pharmacologic probes of lymphocyte signal transduction. *Annu. Rev. Immunol.* **10**, 519–560.
- Sigal N. H., Dumont F., Durette P., Siekierka J. J., Peterson L., Rich D. H., Dunlap B. E., Staruch M. J., Melino M. R., Koprak S. L., Williams D., Witzel B., and Pisano J. M. (1991) Is cyclophilin involved in the immunosuppressive and nephrotoxic mechanism of action of cyclosporin A? *J. Exp. Med.* **173**, 619–628.
- Skene J. H. P. (1990) GAP-43 as a 'calmodulin sponge' and some implications for calcium signaling in axon terminals. *Neurosci. Res.* **13**, S112–S125.
- Skene J. H. P. (1989) Axonal growth-associated proteins. *Annu. Rev. Neurosci.* **12**, 127–156.
- Skene J. H. P. and Willard M. B. (1981a) Changes in axonally transported proteins during axon regeneration in toad retinal ganglion cells. *J. Cell Biol.* **89**, 86–95.
- Skene J. H. P. and Willard M. B. (1981b) Axonally transported proteins associated with growth in rabbit central and peripheral nervous system. *J. Cell Biol.* **89**, 96–103.
- Snyder S. H. and Sabatini D. M. (1995) Immunophilins and the nervous system. *Nature Medicine* **1**, 32–37.
- Sporel-Özkat R. E., Edwards P. M., Gerritsen van der Hoop G., and Gispen W. H. (1990) An ACTH(4–9) analog, Org 2766, improves recovery from acrylamide neuropathy in rats. *Eur. J. Pharmacol.* **186**, 181–187.
- Starr R., Hall F. L., and Moneiro M. J. (1996) A cdc2-like kinase distinct from cdk5 is associated with neurofilaments. *J. Cell Sci.* **109**, 1565–1573.

- Starzl T. E., Todo S., Fung J., Demetris A. J., Venkataramman R., and Jain A. (1989) FK-506 for liver, kidney, and pancreas transplantation. *Lancet* **2**, 1000–1004.
- Starzl T. E., Makowka L., and Todo S. (1987) FK-506: a potential breakthrough in immunosuppression. *Transplant. Proc.* **19**, S3–S104.
- Steiner J. P., Connolly M. A., Valentine H. L., Hamilton G. S., Dawson T. M., Hester L., and Snyder S. H. (1997a) Neurotrophic actions of non-immunosuppressive ligands of immunophilins. *Nature Med.* **3**, 1–8.
- Steiner J. P., Hamilton G. S., Ross D. T., Valentine H. L., Guo H., Connolly M. A., Liang S., Ramsey C., Li J., Huang W., Howorth P., Soni R., Fuller M., Sauer H., Nowotnik A. C., and Suzdak P. D. (1997b) Neurotrophic immunophilin ligands stimulate structural and functional recovery in neurodegenerative animal models. *Proc. Natl. Acad. Sci. USA* **94**, 2019–2024.
- Steiner J. P., Dawson T. M., Fotuhi M., and Snyder S. H. (1996) Immunophilin regulation of neurotransmitter release. *Mol. Med.* **96**, 1076–1551.
- Steiner J. P., Dawson T. M., Fotuhi M., Glatt C. E., Snowman A. M. M., Cohen N., and Snyder S. H. (1992) High brain densities of the immunophilin FKBP colocalized with calcineurin. *Nature* **358**, 584–587.
- Steiner J. P., Dawson T. M., Fotuhi C., Glatt C., Snowman A. M., Cohen N., and Snyder S. H. (1991) The immunosuppressant FK-506 enhances phosphorylation of GAP43: implications for a role in modulation of growth cone function and neurotransmitter release. *Soc. Neurosci. Abstr.* **18**, 603.
- Strand F. L. and Smith C. M. (1986) LPH, ACTH, MSH and motor systems, in *Neuropeptides and Behavior* (de Wied D., Gispen W. H., and van Wimersma Greidanus T. B., eds.), Pergamon, New York, pp. 245–272.
- Strand F. L. and Kung T. T. (1980) ACTH accelerates recovery of neuromuscular function following crushing of peripheral nerve. *Peptides* **1**, 135–138.
- Sugawara T., Itoh Y., Mori E., Kojima H., and Kowada M. (1995) Immunosuppressive agents enhance regeneration of adult dorsal root axons into spinal cord. *Soc. Neurosci. Abstr.* **21**, 315.
- Tai P.-K., Albers M. W., Chang H., Faber L. E., and Schreiber S. L. (1992) Association of a 59-kilodalton immunophilin with the glucocorticoid receptor complex. *Science* **256**, 1315–1318.
- Takei K., Jay D. G., and Mikoshiba K. (1996) Regional laser inactivation of type 1 inositol 1,4,5 trisphosphate receptor affects growth cone morphology and motility in chick DRG neurons. *Soc. Neurosci. Abstr.* **22**, 740.
- Taniuchi M., Clark H. B., Schweitzer J. B., and Johnson E. M., Jr. (1988) Expression of nerve growth factor receptors by Schwann cells of axotomized peripheral nerves: ultrastructural location, suppression by axonal contact, and binding properties. *J. Neurosci.* **8**, 664–681.
- Terada N., Or R., Weinberg K., Domenico J., Lucas J. J., and Gelfand E. W. (1992) Transcription of IL-4 genes is not inhibited by cyclosporin A in competent T cells. *J. Biol. Chem.* **267**, 21,207–21,210.
- Timerman A. P., Ogunbumni E., Freund E., Wiederrecht G., Marks A. R., and Fleischer S. (1993) The calcium release channel of sarcoplasmic reticulum is modulated by FK-506-binding protein. *J. Biol. Chem.* **268**, 22,992–22,999.
- Tocci M. J., Matkovich D. A., Matkovich K., Collier K. A., Kwok P., Dumont F., Lin S., Degudicibus S., Siekerka J. J., Chin J., and Hutchinson N. I. (1989) The immunosuppressant FK506 selectively inhibits expression of early T cell activation genes. *J. Immunol.* **143**, 718–726.
- Todo S., Demetris A., Ueda Y., Imventarza O., Cadoff E., Zeevi A., and Starzl T. E. (1989) Renal transplantation in baboons under FK 506. *Surgery* **106**, 444–451.
- Todo S., Ueda Y., Demetris J. A., Imventarza O., Nalesnik M., Venkataramman R., Makowka L., and Starzl T. E. (1988) Immunosuppression of canine, monkey, and baboon allografts by FK 506: with special reference to synergism with other drugs and to tolerance induction. *Surgery* **104**, 239–249.
- Todo S., Podesta L., ChapChap P., Kahn D., Pan E.-E., Ueda Y., Okuda K., Imventarza O., Casavilla A., Demetris A. J., Makowka L., and Starzl T. E. (1987) Orthotopic liver transplantation in dogs receiving FK-506. *Transplant. Proc.* **19**, S64–S67.
- Tokime T., Nozaki K., and Kikuchi H. (1996) Neuroprotective effect of FK506, an immunosuppressant, on transient global ischemia in gerbil. *Neurosci. Lett.* **206**, 81–84.
- Tong G., Shepherd D., and Jahr C. E. (1995) Synaptic desensitization of NMDA receptors by calcineurin. *Science* **267**, 1510–1512.
- Torii N., Kamishita T., Otsu Y., and Tsumoto T. (1995) An inhibitor for calcineurin, FK506, blocks

- induction of long-term depression in rat visual cortex. *Neuroscience* **185**, 1–4.
- Vincenti F., Laskow D. A., Neylan J. F., Mendez R., and Matas A. J. (1996) One-year follow-up of an open-label trial of FK506 for primary kidney transplantation. *Transplantation* **61**, 1576–1581.
- Walsh C. T., Zydowsky L. D., and McKeon F. D. (1992) Cyclosporin A, the cyclophilin class of peptidylprolyl isomerases, and blockade of T cell signal transduction. *J. Biol. Chem.* **267**, 13,115–13,118.
- Wang M. S., Zeleny-Pooley M., and Gold B. G. (1997) Comparative dose-dependency study of FK506 and cyclosporin A on the rate of axonal regeneration in rat sciatic nerve. *J. Pharmacol. Exp. Ther.* **282**, 1084–1093.
- Wang T., Li B.-y., Danielson P. D., Rockwell S., Lechleider R. J., Martin J., Manganaro T., and Donahoe P. K. (1996) The immunophilin FKBP12 functions as a common inhibitor of the TGF $\beta$  family type I receptors. *Cell* **86**, 435–444.
- Wang T., Donahoe P. K., and Zervos A. S. (1994) Specific interaction of type I receptors of the TGF- $\beta$  family with the immunophilin FKBP-12. *Science* **265**, 674–676.
- Wayne D. B. and Skene J. H. P. (1995) Endogenous nitric oxide modulates neurite growth in regenerating rat sensory neurons. *Soc. Neurosci. Abstr.* **21**, 1292.
- Wiederrecht G. and Etzkorn F. (1995) The immunophilins. *Perspectives Drug Discovery Design* **2**, 57–84.
- Wiederrecht G., Lam E., Hung S., Martin M., and Sigal N. (1993) The mechanism of action of FK-506 and cyclosporin A. *Ann. NY Acad. Sci.* **696**, 9–19.
- Wiederrecht G., Hung S., Chan H. K., Marcy A., Martin M., Calaycay J., Boulton D., Sigal N., Kincaid R. L., and Siekierka J. J. (1992) Characterization of high molecular weight FK-506 binding activities reveals a novel FK-506-binding protein as well as a protein complex. *J. Biol. Chem.* **267**, 21,753–21,760.
- Wijdicks E. F. M., Weisner R. H., Dahlke L. J., and Krom R. A. F. (1994) FK506-induced neurotoxicity in liver transplantation. *Ann. Neurol.* **35**, 498–501.
- Yagita Y., Kitagawa K., Matsushita K., Taguchi A., Mabuchi T., Ohtsuki T., Yanagihara T., and Matsumoto M. (1996) Effect of immunosuppressant FK506 on ischemia-induced degeneration of hippocampal neurons in gerbils. *Life Sci.* **59**, 1643–1650.
- Yao G. L. and Kiyama H. (1995) Dexamethasone enhances level of GAP-43 mRNA after nerve injury and facilitates re-projection of the hypoglossal nerve. *Mol. Brain Res.* **32**, 308–312.

STIC-ILL

*MD*  
*DH301.F4*

**From:** Holleran, Anne  
**Sent:** Friday, December 05, 2003 1:39 PM  
**To:** STIC-ILL  
**Subject:** refs. for 09/780,953

Please send copies of the following papers:

1. Gold Molecular Neurobiology (1997) 15(3): 285-306
2. Jain Transplant Intl. (2002) 15(12): 630-634
3. Piek Faseb J. (1999) 13(15): 2105-2124

Anne Holleran  
AU: 1642  
Tel: 308-8892  
RM: 8e03

mailbox: 8e12

# Specificity, diversity, and regulation in TGF- $\beta$ superfamily signaling

ESTER PIEK, CARL-HENRIK HELDIN, AND PETER TEN DIJKE<sup>1</sup>

Ludwig Institute for Cancer Research, Box 595, S-75124 Uppsala, Sweden

**ABSTRACT** Transforming growth factor- $\beta$  (TGF- $\beta$ ) superfamily members are multifunctional cell-cell signaling proteins that play pivotal roles in tissue homeostasis and development of multicellular animals. They mediate their pleiotropic effects from membrane to nucleus through distinct combinations of type I and type II serine/threonine kinase receptors and their downstream effectors, known as Smad proteins. Certain Smads, termed receptor-regulated Smads, become phosphorylated by activated type I receptors and form heteromeric complexes with a common-partner Smad4, which translocates into the nucleus to control gene transcription. In addition to these signal transducing Smads, inhibitory Smads have been identified that inhibit the activation of receptor-regulated Smads. In contrast to the still growing TGF- $\beta$  superfamily (with ~30 members in mammals), relatively few type I and type II receptors as well as Smads have been identified. We will focus on recent insights into the molecular mechanisms by which signaling specificity between different TGF- $\beta$  superfamily members is achieved and regulated, and how a single family member can elicit a broad scale of biological responses.—Piek, E., Heldin, C.-H., ten Dijke, P. Specificity, diversity, and regulation in TGF- $\beta$  superfamily signaling. *FASEB J.* 13, 2105–2124 (1999)

**Key Words:** activin • bone morphogenetic protein • signal transduction • Smad • transforming growth factor- $\beta$

## THE TGF- $\beta$ SUPERFAMILY AND ITS BINDING PROTEINS

### TGF- $\beta$ superfamily: multiple factors with pleiotropic functions

THE TRANSFORMING GROWTH FACTOR- $\beta$  (TGF- $\beta$ ) SUPERFAMILY is composed of many multifunctional cytokines including TGF- $\beta$ s, activins, inhibins, anti-müllerian hormone (AMH), bone morphogenetic proteins (BMPs), myostatin, and others. The highly similar TGF- $\beta$  isoforms TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 potently inhibit cellular proliferation of many cell types, including those from epithelial origin. Most mesenchymal cells, however, are stimulated in their growth by TGF- $\beta$ . In addition, TGF- $\beta$ s strongly induce extracellular ma-

trix synthesis and integrin expression, and modulate immune responses (reviewed in refs 1, 2). BMPs are potent inducers of bone and cartilage formation and play important developmental roles in the induction of ventral mesoderm, differentiation of neural tissue, and organogenesis (reviewed in refs 3, 4). Activins, named after their initial identification as activators of follicle-stimulating hormone (FSH) secretion from pituitary glands, are also known to promote erythropoiesis, mediate dorsal mesoderm induction, and contribute to survival of nerve cells (reviewed in ref 5). Several growth factors belonging to the TGF- $\beta$  superfamily play important roles in embryonic patterning and tissue homeostasis. Their inappropriate functioning has been implicated in several pathological situations like fibrosis, rheumatoid arthritis, and carcinogenesis.

### Distinct *in vivo* expression patterns of TGF- $\beta$ isoforms

TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 are highly similar in their biological activities *in vitro*. However, they differ in their *in vivo* expression patterns (reviewed in ref 6), which largely explains the unique isoform-specific phenotypes displayed by the TGF- $\beta$  knockout mice. TGF- $\beta$ 1-deficient mice that are born alive undergo early postnatal death due to excessive infiltration of inflammatory lymphocytes and macrophages into several organs (7–9). Half of the mice lacking TGF- $\beta$ 1 die *in utero* due to defects in vasculogenesis and hematopoiesis (10). It is thought that maternally supplied TGF- $\beta$ 1, in combination with redundant expression of TGF- $\beta$  isoforms, contributes to normal development of the embryos. TGF- $\beta$ 1 null mice born from TGF- $\beta$ 1-deficient mothers display abnormal cardiac development (11). TGF- $\beta$ 2 knockout mice show multiple developmental malformations of tissues and organs, leading to perinatal death (12). TGF- $\beta$ 3 null mice die shortly after birth, displaying delayed pulmonary development and defective palate development (13, 14).

<sup>1</sup> Correspondence and current address: Division of Cellular Biochemistry, The Netherlands Cancer Institute, Plesmanlaan 121, 1066 CX Amsterdam, The Netherlands. E-mail: ptdijke@nki.nl




Analysis of the genes encoding the three TGF- $\beta$ s revealed that each isoform is controlled by differentially regulated promoters (reviewed in ref 15). The existence of different genes encoding functionally similar proteins, yet controlled by differentially regulated promoters, might provide an important mechanism to ensure tissue-specific and spatio-temporal expression patterns of the different TGF- $\beta$  isoforms, thereby resulting in proper embryonic development and tissue homeostasis.

In addition, posttranscriptional control mechanisms contribute to regulation of the production of TGF- $\beta$ . TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 that have long, GC-rich 5'-untranslated regions and intramolecular duplex loops located in close proximity to the transcriptional start site negatively modulate TGF- $\beta$  expression, possibly by binding cell type-specific cytoplasmic molecules (16).

### Control of TGF- $\beta$ bioactivity

Members of the TGF- $\beta$  superfamily are synthesized as large precursor molecules that are proteolytically processed in the Golgi apparatus by the convertase family of endoproteases, of which furin is a member. Furin cleaves the precursor into a mature TGF- $\beta$  and amino-terminal precursor remnant, also termed latency-associated protein, or LAP (reviewed in ref 17 and references therein; 18). LAP remains noncovalently linked to TGF- $\beta$  and prevents binding of mature TGF- $\beta$  to its receptors. These so-called small latent TGF- $\beta$  complexes are significantly more stable than bioactive TGF- $\beta$ . Within the Golgi, LAP covalently interacts with latent TGF- $\beta$  binding proteins (LTBPs), to form large, latent complexes (reviewed in ref 19 and references therein). Four different LTBP homologs, LTBPs 1-4, have been characterized, including several alternative splice variants (reviewed in refs 19, 20). LTBPs function to enhance secretion and stability of the TGF- $\beta$ -LAP complex, ensure correct folding of TGF- $\beta$ , and target the latent TGF- $\beta$  complex to the extracellular matrix of certain cells and tissues for storage or to the cell surface where activation takes place (reviewed in refs 17, 19 and references therein). LTBP-1, -2, and -4 contain RGD sequences that are integrin binding sites (20, 21). It has been shown that large latent TGF- $\beta$  complexes directly interact with integrin  $\alpha_v\beta_1$  at the cell surface (21), which may enable TGF- $\beta$  to activate integrin signaling. Activation of latent TGF- $\beta$  into biologically active mature TGF- $\beta$  is controlled by proteases like plasmin or cathepsin, which cleave LAP (Fig. 1), and by binding of LAP to mannose-6-phosphate receptors (17, 19 and references therein). Lipoprotein Lp(a) inhibits activation of plasminogen to plasmin and thereby negatively affects activation of TGF- $\beta$  (reviewed in ref 22 and



**Figure 1.** Activation of latent TGF- $\beta$ . TGF- $\beta$  is synthesized as large latent complexes. In platelets the TGF- $\beta$  latent complex consists of mature TGF- $\beta$  that is noncovalently associated with a disulfide-bonded complex of a dimer of the amino-terminal propeptide of the TGF- $\beta$  precursor (also termed latency-associated peptide, LAP) and a third component, termed latent TGF- $\beta$  binding protein (LTBP). Active TGF- $\beta$  (capable of receptor binding) can be released from the latent complex by specific proteases, like plasmin. This process is likely to occur at the plasma membrane. Lipoprotein Lp(a) is structurally related to plasminogen, the precursor of plasmin. Lp(a) can inhibit plasmin generation, and inhibit activation of latent TGF- $\beta$ . An alternative mechanism for activation of latent TGF- $\beta$  is through thrombospondin, which appears to induce a conformational change in LAP (not shown).

references therein). Other mechanisms that have been implicated in activation of latent TGF- $\beta$  are deglycosylation of LAP (23), exposure to reactive oxygen species (24), or acidic cellular microenvironments (25). In addition, an important activator of TGF- $\beta$  *in vivo* appears to be thrombospondin-1, which induces a conformational change of LAP and thereby activates TGF- $\beta$  (26).

### Signaling diversity by generation of heterodimeric ligands

After proteolytic cleavage of the mature carboxyl-terminal parts, biologically active TGF- $\beta$ 1, - $\beta$ 2, or - $\beta$ 3 homodimers are generated. Homodimers are most abundant, but TGF- $\beta$ 1.2 (27, 28) and TGF- $\beta$ 2.3 (28) heterodimers have been identified *in vivo*. For activation, four different  $\beta$  chains have been identified— $\beta$ A,  $\beta$ B,  $\beta$ C, and  $\beta$ E (reviewed in ref 29 and references therein)—of which  $\beta$ A and  $\beta$ B are known to form homo- as well as heterodimers. Whether heterodimers can also be formed with the  $\beta$ C and  $\beta$ E chains and whether these different isoforms exert different biological activities is not clear. Inhibins, which are heterodimers of inhibin  $\alpha$  chains and activin  $\beta$  chains, are functional antagonists of activin signaling (reviewed in ref 5). In the case of BMPs, BMP2/7 and BMP4/7 heterodimers are much more potent in the induction of ventral mesoderm (30), as well as in bone induction (31) than their respective homodimers. The coexpression of individual BMPs in several tissues suggests that heterodimer formation might occur *in vivo*, and heterodimers of BMP2 and BMP-7 have been isolated from bovine bone (32). The mechanism underlying potentiated signaling by heterodimers compared to homodimers has not been investigated, but might be due to formation



TABLE 1. *TGF- $\beta$  superfamily binding proteins*

TGF- $\beta$ superfamily binding protein	Interacting ligands	Molecular characteristics	Functional properties	Reference
Decorin	TGF- $\beta$ s	Extracellular, proteoglycan	Storage in extracellular matrix, inhibitor of TGF- $\beta$ activity	33
Biglycan	TGF- $\beta$ s	Extracellular, proteoglycan	Storage in extracellular matrix, inhibitor of TGF- $\beta$ activity	33, 34
60 kDa protein	TGF- $\beta$ s	Extracellular, matrix-associated	Inhibitor of ligand-receptor interaction	62
$\alpha$ 2-Macroglobulin	TGF- $\beta$ s, activins	Serum component	Clearance factor	35, 36
Follistatin	Activins, BMPs	Extracellular, heparan-sulfate proteoglycan binding protein	Inhibitor of ligand-receptor interaction	Reviewed in ref 36
Noggin	BMPs	Extracellular, soluble	Inhibitor of ligand-receptor interaction	Reviewed in ref 40
Chordin	BMPs	Extracellular, soluble	Inhibitor of ligand-receptor interaction	Reviewed in ref 40
Cerberus	BMPs, activins	Extracellular, soluble	Inhibitor of ligand-receptor interaction	42
Dan	BMPs	Extracellular, soluble	Inhibitor of ligand-receptor interaction	42
Gremlin	BMPs	Extracellular, soluble	Inhibitor of ligand-receptor interaction	42
TBR-III	TGF- $\beta$ s	Transmembrane, proteoglycan	Accessory receptor, ligand presentation	46, 49, 51
Endoglin	TGF- $\beta$ 1, TGF- $\beta$ 3, activins, BMPs	Transmembrane, proteoglycan	Unknown, signal modulator?	47, 53
Xnr3	BMPs	Intracellular, nodal-related	Inhibitor of ligand synthesis	Reviewed in ref 40

of receptor complexes consisting of, for example, two different type II and two different type I receptors, which may activate downstream signaling in a more potent manner.

#### TGF- $\beta$ superfamily binding proteins


After production of bioactive TGF- $\beta$  superfamily members, several extracellular proteins can bind and modify their activity (Table 1). The extracellular matrix proteoglycans decorin and biglycan have been described to inhibit TGF- $\beta$  activity (33). Transgenic mice lacking biglycan expression show dramatically decreased bone mass, suggesting an important role for biglycan in the constitution of bone, possibly by its function as a source for storage of TGF- $\beta$  superfamily members (34).

The serum protein  $\alpha$ 2-macroglobulin associates with circulating TGF- $\beta$  superfamily members; via its interaction with the  $\alpha$ 2-macroglobulin receptor, it takes care of clearance of the growth factors from serum (35, 36).

Fibronectin can bind TGF- $\beta$  and promote its bioactivity, possibly by changing it from a latent to a bioactive conformation (37), and collagen IV may store bioactive TGF- $\beta$  in the extracellular matrix (38).

Factors known to play important roles during embryonic development are follistatin, noggin, chordin, and Short gastrulation (Sog), the *Drosophila* homologue of chordin (39). These proteins are secreted by the Spemann organizer and compete with BMPs to exert opposing effects on developmental fates of mesoderm and ectoderm (reviewed in ref 40 and references therein). Mice deficient in noggin expression display excessive cartilage formation and failure of joint specification during skeletogenesis (41). The TGF- $\beta$  superfamily member Xnr3 has been reported to interfere with BMP signaling in the *Xenopus* Spemann organizer via an intracellular antagonistic mechanism that prevents BMP synthesis (reviewed in ref 40).

Cerberus, DAN, and gremlin have recently been identified as antagonists of BMP signaling during



**Figure 2.** Mechanism of TGF- $\beta$  receptor activation. Schematic stepwise illustration of the current model for TGF- $\beta$  receptor activation. *A*) TGF- $\beta$ 1 initially binds the accessory TGF- $\beta$  type III receptor (TBR-III), which presents the ligand to TGF- $\beta$  type II receptor (TBR-II). *B*) Subsequently, TGF- $\beta$  bound to TBR-II, recruits TGF- $\beta$  type I receptor (TBR-I) into the complex, thereby forming a heteromeric complex of two TBR-IIs and two TBR-Is. *C*) The constitutively active TBR-II kinase phosphorylates TBR-I. *D*) TBR-I propagates the signal downstream through phosphorylation of particular R-Smads, i.e., Smad2 and Smad3. Recruitment of Smad2 and Smad3 to the TGF- $\beta$  receptor complex is achieved through the Smad anchor for activation (SARA). SARA is membrane associated and capable of binding both R-Smad and the TGF- $\beta$  receptor complex.

early embryogenesis (42). Cerberus can also counteract signaling by activin and nodal. These proteins, like noggin and chordin, have been shown to interact directly with the growth factors, thereby preventing interaction of the ligands with their signaling receptors (reviewed in refs 40, 42). Follistatin counteracts activin signaling by a similar mechanism (reviewed in ref 36), whereas the mechanism by which follistatin interferes with BMP signaling is unclear. BMP-follistatin complexes were found to interact normally with BMP receptors (43).

Interaction of chordin or Sog with BMPs is controlled by metalloproteases of the astacin family, to which tolloid, the *Drosophila* homologue of BMP1, *Xenopus* xolloid, and zebrafish Ztd, belongs, and they are able to proteolytically cleave chordin and Sog, thereby liberating bioactive BMPs (44; reviewed in ref 45).

An extensive number of membrane-bound or transmembrane TGF- $\beta$  binding proteins are known to interfere with TGF- $\beta$  action. The type I and type II serine/threonine kinase receptors, which will be discussed in detail below, are directly involved in TGF- $\beta$  signal transduction. TGF- $\beta$  receptor-III (TBR-III; also known as betaglycan) and endoglin are transmembrane proteins that share high sequence homology in their short cytoplasmic tails rich in serine residues (46, 47). Betaglycan has several high-affinity binding sites for TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 (48) and facilitates binding of TGF- $\beta$  to their type II receptors, a property that is especially important for TGF- $\beta$ 2, which has low intrinsic affinity for TBR-II (Fig. 2) (49, 50). In contrast, soluble betaglycan acts as an antagonist of TGF- $\beta$  bioactivity by preventing the interaction of TGF- $\beta$  with the signaling receptors (46, 51).

The role of endoglin in modulation of TGF- $\beta$  signaling appears to be different from betaglycan. In contrast to betaglycan, endoglin can interact with TGF- $\beta$ 1, TGF- $\beta$ 3, activin A, BMP-2, and BMP-7 and

requires appropriate type I or type II receptors for efficient ligand binding (52). Ectopic expression of endoglin in monocytes results in a selective abrogation of TGF- $\beta$ 1-induced growth inhibition and fibronectin synthesis (53). Furthermore, mutations in the genes encoding endoglin or ALK-1, a putative TGF- $\beta$  type I receptor in endothelial cells, form the basis for the vascular disorder hereditary hemorrhagic telangiectasia type 1 and type 2, respectively (54, 55), suggesting that endoglin and ALK-1 act in a common signal transduction pathway.

In many vascular endothelial cells and hematopoietic progenitor cells, TGF- $\beta$ 1 and TGF- $\beta$ 3 are equipotent in mediating signaling, but TGF- $\beta$ 2 has weaker biological effects (47, 56). This difference in potency between different TGF- $\beta$  isoforms can be accounted for by the lack of betaglycan in these cells. Differential activities of TGF- $\beta$  isoforms in other cell types have been observed, but their modes of action have not been correlated to receptor binding properties (57, 58). Several cell membrane-associated TGF- $\beta$  binding proteins, including certain glycosyl-phosphatidylinositol (GPI)-anchored proteins (59, 60), have been identified that are endowed with specific affinity for certain TGF- $\beta$  isoforms, but their role in TGF- $\beta$  signaling is not clear. Furthermore, a 60 kDa TGF- $\beta$  binding protein has been characterized (61, 62) that exhibits TGF- $\beta$ 1 binding specificity and interferes with binding of TGF- $\beta$  to the signaling receptors (62).

## SIGNALING VIA TGF- $\beta$ SUPERFAMILY RECEPTORS

**TGF- $\beta$  superfamily members induce formation of heteromeric complexes of type I and type II receptors**

Type I and type II serine/threonine kinase receptors are directly involved in signaling by TGF- $\beta$  superfamily

ily members (Fig. 2). TGF- $\beta$ 1, TGF- $\beta$ 3, and activins initially bind to their corresponding type II receptors, after which the type I receptors are recruited into the signaling complex. Type I and type II receptors have an intrinsic affinity for each other, which contributes to stability of the heteromeric complex (63). TGF- $\beta$ 2, which does not bind to T $\beta$ R-II or T $\beta$ R-I alone, can interact with and stabilize transiently formed complexes of T $\beta$ R-I and T $\beta$ R-II (64). BMPs have low affinity for type II or type I receptors individually, and high-affinity binding requires formation of a heteromeric type I/type II receptor complex (65, 66). At present, only a few type I and type II receptors for the TGF- $\beta$  superfamily are known. However, depending on the ligand, multiple heterotetrameric complexes consisting of two type I and two type II receptor homodimers are possible (67). In view of the existence of heterodimeric ligands, it is conceivable that two different type I and two different type II receptors constitute the receptor complex, thereby creating combinatorial signaling.

TGF- $\beta$ s form heteromeric complexes between T $\beta$ R-II and T $\beta$ R-I in most cell types (68, 69). In endothelial cells TGF- $\beta$  can bind and signal through activin receptor-like kinase 1 (ALK-1); ALK-1 has therefore been implicated as an endothelial-specific TGF- $\beta$  type I receptor (T. Imamura, P. ten Dijke, and K. Miyazono, personal communication). Activins signal via combinations of ActR-II or ActR-IIB and ActR-IB (68, 70). Activins have also been shown to interact with ActR-I (68, 71), although this receptor appears to play a minor role in activin signaling. BMPs interact with ActR-II, ActR-IIB, or BMPR-II, in combination with ActR-I, BMPR-IA, or BMPR-IB (65, 66, 72). For AMH, only a type II receptor has been identified (73). A potential AMH type I receptor is ActR-I, which is coexpressed with AMHR-II in the Mullerian duct and can mediate AMH-specific repression of FSH-induced aromatase activity (A. Themmen, personal communication). ALK-7 is an orphan receptor (74). A novel type I receptor, designated ZALK-8, was recently cloned and shown to be widely expressed throughout early zebrafish development (75). Mammalian homologues of ZALK-8 as well as of zebrafish TARAM-A, which is a type I receptor involved in induction of anterior dorsal mesoderm (76), have not yet been identified. The ligands for these receptors have not been elucidated.

#### Activation and regulation of serine/threonine kinase type I and type II receptors

Type I receptor kinases become activated upon phosphorylation by the constitutively active type II receptor kinase (Fig. 2). Thus, type I receptors act

downstream of type II receptors (77). Consistent with this finding, type I receptors were found to determine the specificity of the intracellular signals induced by different TGF- $\beta$  superfamily members, whereas the L45 region in type I receptors was found to be important in determining the specificity of type I receptors (78–80).

The phosphorylation status of T $\beta$ R-II has been shown to influence TGF- $\beta$  signaling. T $\beta$ R-II is a constitutively active kinase that requires phosphorylation of Ser213 (located outside the kinase domain) to mediate its autocatalytic effect. Ligand-dependent autophosphorylation of T $\beta$ R-II on either Ser409 or Ser416 differentially contributes to regulation of its activity, leading to stimulation or inhibition of TGF- $\beta$  signaling, respectively (81).

Differential kinetics in biosynthesis, ligand-induced internalization, and down-regulation of type I vs. type II receptors have been attributed to differential modulation of TGF- $\beta$  signaling (82–85). The half-life and processing of T $\beta$ R-II in the endoplasmic reticulum are considerably shorter than that of T $\beta$ R-I (82–85). Furthermore, homo- and heteromeric receptor complexes have distinct endocytotic fates (84). Rapid modulation of T $\beta$ R-II expression levels as well as ligand-bound T $\beta$ R-I/T $\beta$ R-II complexes may be important for fine-tuning of signaling by TGF- $\beta$ .

Diversity in receptor availability might be provided by the existence of alternative receptor transcripts, as identified for ActR-IIA (86), ActR-IB (87), ActR-IIB (70), AMHR-II (73), and T $\beta$ R-II (88). In the case of ActR-IIB, differential affinity of activin for the alternatively spliced gene products has been observed. In addition, differences in the cytoplasmic domains offer possibilities for differential modes of signaling (70).

Several growth factors are known to bind to the same type II receptor—for example, activins and BMPs, which share ActR-II and ActR-IIB (72, 89). This enables signaling by several growth factors via a limited scale of available receptor types, whereas it can lead to competition of different ligands for the same type II receptors, thereby fine-tuning signaling via type I receptors. This is the case for activin and inhibin, since inhibin is known to counteract many of activin functions by competing for binding to the activin type II receptor (90). A similar mechanism has been suggested to underlie the inhibitory action of zebrafish activin, most closely related to mouse lefty, upon activin signaling (91), and appears to contribute to the functional antagonism observed between activin and OP-1 in Tera-2 embryonal carcinoma cells (89). Whether different ligands might also compete for binding to common type I receptors is not clear at present.



TABLE 2. *TGF- $\beta$  superfamily receptor interacting proteins*

Receptor-interacting protein	Interacting receptor type	Molecular characteristics	Functional properties	Reference
FKBP12	Type I receptors	Immunophilin	Inhibitor of T $\beta$ R-I kinase	97, 98
FT $\alpha$	TGF- $\beta$ type I receptor	Farnesyl transferase	Unknown	99
TRIP-1	TGF- $\beta$ type II receptor	WD-40 repeat protein	Signal modulator	92, 93
STRAP	TGF- $\beta$ type I, II receptors	WD-40 repeat protein	Inhibitor of transcriptional responses	94
TRAP-1	TGF- $\beta$ type I receptor	Without motif	Inhibitor of TGF- $\beta$ signaling	96
B $\alpha$	TGF- $\beta$ type I receptor, activin type I receptor	WD-40 repeat protein, phosphatase 2A-subunit	Enhancer of growth inhibition	95
Apolipoprotein J	TGF- $\beta$ type I, II receptors	Nuclear protein or secreted glycoprotein	Role in TGF- $\beta$ receptor signaling and/or processing	227
SARA	TGF- $\beta$ receptor complex	FYVE-containing protein	Smad subcellular localizer	147
Smad	Type I receptors	Transcription factor	Signal transducer	29, 101
BRAM-1	BMP type IA receptor	E1A-associated protein	Unknown	212
XIAP	BMP type IA receptor	Inhibitor of apoptosis protein	Adaptor protein for TAB1, inhibitor of apoptosis	211

### Cytoplasmic TGF- $\beta$ receptor interacting proteins distinct from Smads

A number of cytoplasmic proteins are now known to interact with the kinase domain of type I and type II receptors, thereby manipulating or mediating their signaling capacities (Table 2). TRIP-1, a protein that contains five WD domains, is known to bind and become phosphorylated by T $\beta$ R-II. The interaction requires a functional T $\beta$ R-II kinase (92). Overexpression of TRIP-1 attenuates TGF- $\beta$ -induced PAI-1 transcriptional response, but not TGF- $\beta$ -induced cyclin A response (93). The WD domain-containing protein STRAP can associate with T $\beta$ R-I and T $\beta$ R-II. STRAP and the inhibitory Smad7 synergistically block TGF- $\beta$ -mediated transcriptional activation (94). Another WD-40 repeat protein B $\alpha$ , a subunit of protein phosphatase 2A, has been reported to interact with and become phosphorylated by T $\beta$ R-I. B $\alpha$  potentiates the antiproliferative effect mediated by TGF- $\beta$  (95). T $\beta$ R-I-associated protein-1 (TRAP-1) interacts specifically with the activated T $\beta$ R-I. However, its functional role in TGF- $\beta$  signaling is unclear (96).

The immunophilin FKBP12 interacts with type I serine/threonine kinase receptors (97) through an FK506-like Leu-Pro motif preceding the kinase domain (98). FKBP-12 inhibits phosphorylation of the type I receptors by the type II receptors, possibly by sterical hindrance, and may function to prevent leaky signaling in the basal cell state (99). FKBP12 dissociates from T $\beta$ R-I after ligand-induced phosphorylation of T $\beta$ R-I by T $\beta$ R-II (97, 98).

Farnesyltransferase  $\alpha$  (FT $\alpha$ ) has also been shown to interact with and become phosphorylated by activin and TGF- $\beta$  type I receptors (99). However, farnesyl transferase activity was shown to be dispensable for TGF- $\beta$  signaling (100).

### Type I receptors as determinants of downstream signal propagation

Activation of type I receptors by type II receptors occurs by phosphorylation of serine and threonine residues within a juxtamembrane region preceding the kinase domain of type I receptors that is characterized by repeats of glycine and serine residues (also termed the GS domain) (77). Specificity of signal propagation to Smad molecules, the main downstream components in serine/threonine kinase receptor signaling, is determined by type I receptors (reviewed in refs 29, 101). More specifically, a region flanking  $\beta$  strands 4 and 5 in the kinase domain, designated L45 loop, confers a high degree of specificity in Smad interaction with the type I receptors (78, 80). The L45 loop is highly conserved among type I receptors with comparable signaling specificity, but differs significantly between BMP and TGF- $\beta$  or activin type I receptors (79, 80). Swapping the L45 loop between T $\beta$ R-I and BMPR-IB, followed by ligand-stimulation, was shown to result in exchange of Smad1 vs. Smad2 recognition specificity, accompanied by a switch in transcriptional responses (79, 80). However, differential effects between the two BMP type I receptors have been described (102-104). CDMP-1 and CDMP-2, which both interact with BMPR-IB and BMPR-II, differentially regulate osteogenesis (105, 106). Thus, in addition to a role for the L45 loop, other mechanisms must contribute to specify downstream signaling after activation of type I receptors.

It has been shown that the cytoplasmic juxtamembrane region of T $\beta$ R-I is involved in specifying signal transduction, since mutations of certain serine or threonine residues within this region selectively impair TGF- $\beta$ -mediated growth inhibition, but do not affect TGF- $\beta$ -induced PAI-1 or fibronectin synthesis

(107). However, *in vivo* phosphorylation of these sites by T $\beta$ R-II has not yet been reported. Mutation of Ser165, located amino-terminal of the GS domain, results in potentiation of TGF- $\beta$ -mediated growth inhibition and extracellular matrix induction but diminishes TGF- $\beta$ -induced apoptosis (108). Thus, additional residues in T $\beta$ R-I appear to modulate distinct pathways triggered by TGF- $\beta$ .


In certain cell systems it has been observed that the expression level of type I receptors correlates with extracellular matrix production and gene transcription, whereas the expression level of type II receptors correlates to the growth inhibitory response, suggesting participation of type I and type II receptors in different signaling pathways. However, the observations made are all compatible with an alternative interpretation, which is that all signals are initiated by activation of the type I receptor and that different activation thresholds of the signaling pathways leading to growth inhibition vs. gene transcription underlie the apparent differential modes of signaling by type I and type II receptors (109).

## INTRACELLULAR SIGNALING BY TGF- $\beta$ SUPERFAMILY MEMBERS

### Receptor-activated Smads and common-partner Smads as mediators of TGF- $\beta$ signal transduction

Genetic studies in *Drosophila* and *Caenorhabditis elegans* have led to identification of a conserved family of intracellular signal transducers for TGF- $\beta$  superfamily members. The founding member, *Mothers against dpp* (*Mad*) (110, 111) was isolated from a genetic screen for enhancers of a weak *Dpp* maternal phenotype. Subsequently, three *Mad* homologues, called *Sma-2*, *Sma-3*, and *Sma-4*, were picked up in *C. elegans* by searching for genes whose mutations caused the same small body-size phenotype as observed for mutant *Daf-4*, a type II serine/threonine kinase receptor (112). At present, eight different *Sma*- and *Mad*-related proteins have been identified in mammals, and are termed Smads (Fig. 3A). They can be subdivided into three distinct subclasses: receptor-activated Smads (R-Smads), common-partner Smads (Co-Smads), and inhibitory Smads (anti-Smads). R-Smads and Co-Smads are homologous in their amino- and carboxyl-terminals, called the MH1 and MH2 domains, respectively. These domains are connected by a proline-rich linker region. R-Smads contain SSXS phosphorylation motifs in their own carboxy termini.

Smad1, Smad5, and Smad8 are involved in BMP signaling and become phosphorylated by ActR-I, BMPR-IA, or BMPR-IB (113–117). Smad2 and Smad3 are mediators of TGF- $\beta$  and activin signaling



**Figure 3.** The Smad family of intracellular signal transducers for TGF- $\beta$  superfamily proteins. **A)** A phylogenetic analysis of mammalian Smad proteins. **B)** The Mad homology (MH)1 and MH2 domains, have affinity for each other; type I receptor-mediated phosphorylation of R-Smads at their carboxy termini may induce a conformation change, thereby relieving the autoinhibitory effect of MH1 on MH2 and vice versa. **C)** Schematic structure of R-Smad and Co-Smad and their functional domains. Direct DNA binding is mediated through the MH1 domain whereas MH2 has transcriptional activation activity. The sites of phosphorylation in R-Smads by type I receptors at the extreme carboxy terminus and by ERK in linker region are indicated.

and interact with and become phosphorylated by T $\beta$ R-I or ActR-IB (118–121). Consistent with these findings, injection of BMP-Smads into *Xenopus* animal caps results in ventralization of mesoderm (115, 116, 122, 123), and ectopic expression of Smad2 in *Xenopus* embryos induces dorsal mesoderm and secondary axis formation analogous to activin/Vg-1-like responses (122, 123). Smad2 null mice lack anterior-posterior specification and fail to develop mesoderm (124, 125). The phenotype of Smad2 knockouts is strikingly similar to Nodal knockouts and suggests that they cooperate in regulation of gastrulation (124). Mutant mice lacking Smad3 expression have been reported to be viable and fertile; however, Smad3<sup>-/-</sup> animals were found to exhibit limb malformations (126), have a defect in immune function (127), and develop colon carcinomas 4–6 months after birth (128). Smad3 null cell lines showed strongly impaired TGF- $\beta$  responsiveness and indicated an important role for Smad3 in TGF- $\beta$ -mediated inhibition of cellular proliferation (126). The Smad3 knockout phenotype observed by Zhu and co-workers (128) classifies Smad3 as a tumor suppressor gene, although Smad3 mutations have not yet been detected in tumorigenesis (129). TGF- $\beta$ -mediated phosphorylation of Smad1 has

been observed in breast tumor cells, but the receptor by which this phosphorylation is mediated has not been determined (130). Smad5 phosphorylation in response to TGF- $\beta$  was detected in CD34+ cells (131). ALK-1 has been shown to bind TGF- $\beta$  and to phosphorylate and activate BMP-receptor-regulated Smads (113, 114; T. Imamura, P. ten Dijke, and K. Miyazono, personal communication), and thus could possibly mediate TGF- $\beta$ -induced Smad phosphorylation in these cells.

Three Co-Smads have been characterized: mammalian Smad4/DPC4 (deleted in colon carcinoma; ref 132), *C. elegans* Sma-4 (112), and *Drosophila* Medea (133). Co-Smads play a critical role in signaling by TGF- $\beta$  superfamily members, illustrated by the absence of TGF- $\beta$  signaling and R-Smad functioning in Smad4-deficient tumor cell lines, as well as by the synergistic action of Smad4 with R-Smads in transcriptional activation of target genes and induction of mesoderm (121, 134–137). The importance of Smad4 functioning is also shown by its tumor suppressor activity; mutation or deletion of Smad4 is often associated with tumorigenesis (132, 138–141). Cell lines expressing Smad4-mutants found in human cancers failed to transcriptionally activate transfected luciferase reporters containing Smad binding elements (140). Mice lacking expression of Smad4 show a BMP knockout phenotype, characterized by growth retardation and gastrulation defects (142). However, the abnormalities in mesoderm development of Smad4 null mice could be overcome by the presence of wild-type extraembryonic tissue, indicating that Smad4 is not critical for early gastrulation (142). Although several explanations are possible, one reasonable argument is that a yet unidentified Co-Smad is involved in mediating early embryonic differentiation induced by members of the TGF- $\beta$  superfamily. At later stages of development, Smad4 is important for anterior specifications (142).

Additional indications for the possible existence of novel Co-Smads comes from studies in *Drosophila* where Medea, in contrast to Mad, is dispensable for oogenesis (143). An explanation for the fact that Mad knockouts have a more severe phenotype than Medea knockouts might be that Mad is able to signal certain responses independent of Medea. Furthermore, mutations in Medea efficiently block signaling by Saxophone (Sax) but fail to fully suppress signaling via Thick Veins (Tkv), both of which are Dpp type I receptors (133). Although Medea might be abundantly present to allow Tkv signaling even when hypomorphic alleles are expressed, the inability to fully abrogate Tkv signaling might also reside in the fact that another, yet unidentified, Co-Smad is involved in Tkv signaling or that Tkv can signal independent of Medea. Besides the XSmad4 ortholog, another Smad4/Medea-like protein was recently

identified in *Xenopus* (144, 145; C. Hill, personal communication).

### Mechanisms controlling Smad activation

R-Smads occur as monomers in the cytoplasm (146) and, after ligand-induced type I receptor activation, specifically interact with the L45 loop in the kinase domain of type I receptors (79, 80). Recruitment of R-Smads to the TGF- $\beta$  receptors is promoted by the action of SARA, a FYVE zinc finger domain-containing protein that interacts with both TGF- $\beta$  receptors and Smad2 or Smad3 (Fig. 2) (147, 148). The highly conserved L3 loop of R-Smads, which protrudes from the MH2 core domain as observed in the Smad4-MH2 crystal structure (139), is involved in specifying type I interactions with T $\beta$ R-I and BMPR-I (150). This loop varies between Smad1 and Smad2 in only two amino acids, and exchanging these residues between Smad1 and Smad2 enables their binding to T $\beta$ R-I and BMPR-I, respectively (79, 149). The  $\alpha$ -helix 1 of Smad1, located near the L3 loop in the 3 dimensional structure, has been shown to be important for the interaction of Smad1 with ALK-1 (114).


After type I receptor interaction, R-Smads become activated by phosphorylation of their carboxyl-terminal SS(M/V)S motifs (Fig. 3B) (119, 150–152), which triggers their homodimerization and heteromerization with Co-Smads (134, 153) or other R-Smads (121, 146). This leads to the speculation that homomeric R-Smads, heteromeric R-Smads, and R-Smad/Co-Smad complexes might control different biological responses.

In contrast to R-Smads, Co-Smads do not contain the carboxyl-terminal phosphorylation motif SSXS and are not phosphorylated by the activated type I receptor (119, 120). Although the L3 loop of Co-Smads shares high amino acid similarity with the L3 loop of R-Smads, Co-Smads do not interact with type I receptors (120); instead, their L3 loops are important for heteromerization with phosphorylated R-Smads (139). Resolution of the crystal structure of the Smad4-MH2 domain and analyses of Smad-deletion mutants (139) have indicated that the MH2 domains of R-Smads and Co-Smads are involved in formation of homo- and heteromeric trimers. After association of R-Smads with Co-Smads, the heteromeric complexes are translocated into the nucleus (see Fig. 5).

### Domains controlling nuclear translocation

Whereas ligand-induced nuclear translocation of R-Smads is not dependent on expression of Co-Smads (136), Co-Smads require association with activated R-Smads in order to enter the nucleus (136, 151). A carboxyl-terminal truncation of Smad4, correspond-





**Figure 4.** Smad interacting proteins. Overview of the interacting proteins for Smad3 and Smad4. All proteins that interact with Smad3 also interact with Smad2 except for c-Jun, c-Fos, and Evi-1. The interaction of TGF- $\beta$  receptor with Smad2 or Smad3 occurs through the L3 loop in the MH2 domain, and association of FAST-1 with Smad2 or Smad3 is determined by an exposed  $\alpha$ -helix 2 in MH2 domain.

ing to a mutation of Smad4 found in pancreatic tumors (132) and identified in an inactive form of Mad in *Drosophila* (111), dominant-negatively interferes with nuclear transport of all R- and Co-Smads (137). In contrast, a comparable mutation in R-Smads does not affect their nuclear localization (137). This indicates that the carboxyl-terminal part of Co-Smads is required for nuclear trafficking (137), whereas phosphorylation of R-Smads, though required for their functionality, is not obligatory for nuclear transport. How these mutant Co-Smads interfere with the nuclear transport function of R-Smads needs to be further studied. Removal of the NH<sub>2</sub> terminus and linker region of R-Smads results in strong, ligand-independent nuclear localization (122, 137), whereas an additional carboxyl-terminal truncation of the remaining MH2 domain results in cytoplasmic accumulation at the nuclear membrane (137). Presumably, ligand-induced heteromerization results in a conformational change of the Smad molecules, thereby exposing nuclear translocation signals in R- and Co-Smads. Whether the cytoplasmic residence of R-Smads and Co-Smads in the absence of ligand is due merely to autoinhibition of the MH2 effector domain by association with the MH1 domain (154), or whether a cytoplasmic tether exists that anchors R- and Co-Smads in the cytoplasm in the absence of ligand, is not clear (137).

#### R-Smads and Co-Smads as sequence-specific transcriptional activators

Once in the nucleus, R-Smads and Co-Smads are involved in transcriptional regulation of target genes (Fig. 3C) (reviewed in ref 155), and the MH1 and MH2 domains differentially contribute to this property. Direct binding of Smads to DNA was first shown by the interaction of the *Drosophila* Mad-MH1 domain with specific sequences in the promoter of the Dpp target gene *vestigial* (156). A prerequisite for the interaction of Mad, Smad3, but not Smad4, with DNA is the dissociation of the MH2 domain from the MH1 domain, which is achieved by ligand-induced phosphorylation of the SSXS motif in R-Smads (154,

156). Thus, the MH2 domain can exert an autoinhibitory action on the MH1 DNA binding domain.

In several promoters of genes induced by TGF- $\beta$  signaling like *type VII Collagen*, *JunB*, and *PAI-1* (157, 158, 160), unique elements have been identified with which Smad3 and Smad4 physically interact. The sequences of the Smad binding elements (SBEs) in these promoters are highly related and indicate an AGAC-containing motif as binding site for Smad3 and Smad4. This sequence was also picked up in a random screening for Smad-interacting DNA sequences (161), and mutation of these SBEs in the PAI-1 promoter impairs TGF- $\beta$  responsiveness (160).

Crystallization of the Smad3 MH1 domain bound to an optimal Smad binding sequence revealed that the so-called  $\beta$ -hairpin loop, protruding from the MH1 core domain, interacts with DNA (162). The  $\beta$ -hairpin loop is highly conserved among R-Smads and Co-Smads. However, Smad2 contains an extra exon preceding this region, encoding 30 amino acids, that might interfere with the correct conformation of the  $\beta$ -hairpin loop, in contrast to Smad3 and Smad4. Smad2 has not yet been shown to interact with DNA. Removal of exon 3 in Smad2 enables its efficient binding to DNA (163, 164).

In several TGF- $\beta$  target genes, multiple copies of SBEs can be identified, often located in close proximity to sites for other transcription factors (158, 160, 165–167). Smads have been shown to mediate transcriptional activation by interaction with other regulatory promoter units, like the TPA-responsive elements (TREs)/AP-1 sites in the collagenase promoter, which actually overlap in sequence with SBEs (158, 166). Smads cooperate with the transcription factor TFE3 in the regulation of the *PAI-1* gene (168). The p21/WAF1/Cip1 and p15/INK4B gene promoters lack SBEs, but TGF- $\beta$  transcriptionally regulates these genes via the Sp1 site (169–171). It has been suggested that the interaction of Smads with additional transcription factors such as AP-1 (Fig. 4; 166, 172, 173), forkhead activin signal transducers (FAST; 165, 167, 174), or Sp1 (169, 171) confers additional DNA binding specificity to the Smad-containing transcriptional complex (see below).

## **R-Smads and Co-Smads as transcriptional (co-)activators**

Studies with *Xenopus* explants, in which injection of the Smad2-MH2 domain fully induced dorsalization of mesoderm and secondary axis formation analogous to activin signaling (122), suggested an effector function for MH2 domains. Furthermore, transcriptional activation properties of R- and Co-Smads were shown by fusion constructs of Smad1-MH2 or Smad4-MH2 domains with Gal4-DNA binding domains (175). Heteromerization of Co-Smads with R-Smads is obligatory for their transcriptional activation properties (136, 153, 176).

Several domains in Smad4 have now been identified to contribute to its important role in signaling by members of the TGF- $\beta$  superfamily. As mentioned above, the MH1 domain enables the interaction of R-Smad/Co-Smad complexes with DNA, which is of particular importance for Smad2-Smad4 heteromeric complexes (136) since Smad2 is not endowed with DNA binding capacities. The MH1 domain autosuppresses the MH2 effector domain (154), and heteromerization of Co-Smads with phosphorylated R-Smads has been suggested to release the inhibitory association (177). Studies in which chimeras between Smad1 and Smad4 were tested revealed that the proline-rich linker region of Smad4 contains a domain that is important for its transcriptional activity mediated by the MH2 domain (178). This Smad activation domain (SAD) cooperates functionally with the transcription coactivator MSG1 (178, 179). MSG1 strongly enhances Smad4 transcriptional activity, provided that Smad4 is present in a heteromeric complex with R-Smads (179). Smad heteromerization possibly introduces a configuration that allows *trans*-activators to associate and endow Smad4 with transcriptional potencies (179).

## **Smads interact with transcription factors**

The first evidence that Smads interact and cooperate with transcription factors comes from the interaction observed between Smad2, Smad4, and *Xenopus* FAST-1 (XFAST-1), making a complex that interacts with the activin responsive element (ARE) in the *Xenopus* Mix2 promoter (see Figs. 4, 5) (136, 180, 181). Human FAST-1 and mouse FAST-2 (mFAST-2) have been identified as well, but share little sequence homology with XFAST-1 (165, 167, 174). FAST-1 efficiently interacts with DNA, and Smad2 and Smad4 contribute in providing additional DNA interaction properties (136, 167, 174, 180, 181). Phosphorylation of the Smad2 SSMS motif is required for interaction of  $\alpha$ -helix 2, exposed from the MH2-trimer structure, with FAST-1 (136, 167, 180, 181).

Whereas Smad2/Smad4/mFAST-2 complexes drive transcriptional activation of the goosecoid reporter, Smad3/Smad4/mFAST-2 complexes were found to suppress promoter activation, presumably because Smad3 and Smad4 share the same DNA binding site (165). This suggests that the relative expression levels of Smad2 and Smad3 in a cell might determine the outcome of certain biological responses induced by TGF- $\beta$ .

Additional evidence for differential roles of Smad2 and Smad3 comes from observations in human keratinocytes, where TGF- $\beta$  stimulates Smad2 and Smad3 to comparable extents while activin predominantly activates Smad3 (182). In human lung epithelial cells, Smad3 more potently induces apoptosis compared to Smad2 (183).

## **Cooperation of Smads with the AP-1 complex**

Characterization of Smad binding sites in a collagenase promoter revealed that Smads can mediate transcription via 12-O-tetradecanoyl-13-acetate (TPA)-responsive elements (TREs; 120, 121, 158), to which the transcription factor AP-1, a dimer of c-Jun and c-Fos, binds. These AP-1 binding sites overlap in sequence with consensus Smad binding sites (158). Smad3 directly interacts with TRE and Smad3 and Smad4 activate the TGF- $\beta$ -inducible transcription of TRE-Luc reporter in the absence of c-Jun and c-Fos (166). However, TGF- $\beta$ -induced reporter activity was further enhanced in the presence of c-Jun-c-Fos heterodimers (166). After TGF- $\beta$ -induced Smad activation, Smad3 and Smad4 can interact with JunB, c-Jun, and JunD as well as with c-Fos (166, 172). By footprint analysis and electromobility shift assays, it was shown that c-Jun and Smad3 can bind simultaneously to overlapping sequences in the TRE (166). Crystallographic analysis of the Smad3-MH1 domain bound to DNA indicated that a conformational change of the NH<sub>2</sub> terminus of c-Fos would allow the multimeric Smad-AP-1 complex to interact with a single AP-1 site (162). This cross talk might be an important level of coordination between Smads and MAPK/JNK signaling.

## **Interaction with the coactivators p300/CBP**

Insight into how Smads mediate their transcriptional activating function was recently unraveled by the interaction of R-Smads with the coactivators p300 and CREB binding protein (Fig. 4) (CBP; 176, 184-187). These coactivators contribute to transcriptional activation by loosening the chromatin structure or by increasing the affinity of certain transcription factors for DNA through their intrinsic (or associated) acetyltransferase activity. In addition, they act as a bridging factor between transcription factors and the basal transcription machinery (184



and references therein). In view of the importance of Smad4 for transcriptional activation of the Smad-p300/CBP complex, Smad4 might function as a coactivator as well (176). The interaction between the MH2 domain of R-Smads and CBP/p300 is triggered by ligand-induced phosphorylation of the carboxyl-terminal SSXS motif (176, 184, 186). Phosphorylation most likely induces a conformational change in the R-Smad molecule, which then results in exposure of the CBP/p300 binding site; deletion of the NH<sub>2</sub> terminus and part of the carboxy terminus, leaving the CBP binding site intact, enhances the affinity of Smad3 for CBP/p300 (184). The carboxyl-terminal region of CBP is required for Smad interaction (176, 184, 186) and can associate with other transcription factors like the adenoviral oncoprotein E1A. E1A is known to interfere with TGF- $\beta$ -mediated signaling (188, 189) and is shown to abrogate Smad-mediated transcriptional activation (176, 184, 186), most likely by direct competition with Smads for binding to CBP/p300 (188, 189).


### Negative regulation of Smad function

Activin and BMPs have been shown to compete with each other in *Xenopus* mesoderm development due to limiting amounts of Smad4 (190). In view of the limited amounts of CBP/p300 in the cell (184) and the observed competition between different signal transduction pathways at the level of coactivator availability (191), competition between members of the TGF- $\beta$  superfamily as observed in embryogenesis can also be expected to occur at the level of CBP/p300.

Interaction of the first zinc finger domain of the nuclear protein Evi-1 with the MH2 domain of Smad3 interferes with binding of Smad3 to DNA (Fig. 4), thereby repressing transcriptional activity of Smad3 as well as growth inhibition (192). Evi-1 shows a spatial and temporal expression pattern during mouse embryogenesis, suggesting it could play a regulatory role during development (193).

### Inhibitory Smads

Mammalian Smad6, Smad7, and *Drosophila* Dad have been characterized as inhibitors of TGF- $\beta$  signal transduction (Fig. 5) (194–198). Whereas their amino-terminal domains are highly diverse and share only weak similarity to other Smads, they are homologous to the R- and Co-Smads in their MH2 domains. However, inhibitory Smads lack the carboxyl-terminal SSXS phosphorylation motif, which may enable them to stably associate with type I receptors and to interfere with receptor binding and activation of R-Smads. Phosphorylation of the type I receptors by the type II receptors is essential for inhibitory Smad



**Figure 5.** TGF- $\beta$  superfamily signaling through signal-transducing Smad and inhibitory Smad proteins. After type I receptor activation, R-Smads become phosphorylated, form homomeric complexes with each other, and assemble into heteromeric complexes with Co-Smad, Smad4. Trimers are shown, but hexamers have not been excluded. The stoichiometry between the components is unclear. The heteromeric complexes translocate into the nucleus, where they regulate, in combination with other transcription factors, transcription of target genes. Smad2, Smad4, and FAST-1 are components of an activin-responsive factor that interacts directly in an activin-dependent manner with an activin response element of the *Mix-2* promoter. Inhibitory Smads act opposite from R-Smads by competing with them for interaction with activated type I receptors or by directly competing with R-Smads for heteromeric complex formation with Co-Smad. Smad7 appears to be a general inhibitor of TGF- $\beta$  superfamily signaling whereas Smad6 preferentially inhibits BMP-induced responses.

association (194, 196, 199) and the MH2 domain of anti-Smads suffices in exhibiting the inhibitory effect (194, 196, 199, 200), as manifested by inhibition of R-Smad phosphorylation and abolition of their heteromerization with Co-Smad and nuclear translocation, as well as abrogation of growth factor-induced transcriptional responses and growth inhibition (194–196, 198–202). It has been reported that Smad6 can exert its inhibitory effect via an alternative mode that involves BMP-induced association of Smad6 with phosphorylated Smad1, thereby competing with activated Smad1 for heteromerization with Smad4 (200) (Fig. 5).

The expression of the inhibitory Smads is quickly induced upon stimulation by TGF- $\beta$  superfamily members and thereby provides an autoinhibitory mechanism in TGF- $\beta$  signaling (196, 197, 201, 203). The human Smad7 promoter contains an SBE consensus site that, when fused to a luciferase gene, induces luciferase activity after TGF- $\beta$  stimulation (G. Brodin and R. Heuchel, personal communication). Thus, after receptor activation, R- and Co-

Smads might be directly involved in transcriptional regulation of the Smad7 gene.

The MH1 domain of inhibitory Smads might play a role in specification of their inhibitory actions: *Xenopus* and mouse Smad7, which share 96% identity in their carboxyl-terminal domain but only 51% in the amino-terminal region, exert differential effects in inhibition of TGF- $\beta$  and activin signaling (199). Mouse Smad7 inhibits T $\beta$ R-I-induced phosphorylation of Smad2 and Smad3 (194, 196, 199), BMPR-IA and BMPR-IB-mediated phosphorylation of Smad1 and Smad5 (199), and ActR-I mediated phosphorylation of Smad1 (199); it therefore appears to be a common inhibitor of R-Smad activation. Whereas Smad6 has been implicated in inhibition of TGF- $\beta$  signaling (194, 204), it appears to play a more pronounced role in inhibition of BMP signaling (Fig. 5) (195, 200, 201, 203, 205, 206). The mechanism of action for inhibitory Smads in TGF- $\beta$  superfamily signaling is evolutionarily well conserved; it has recently been shown that Dad, which is a Dpp-inducible inhibitory Smad in *Drosophila* (197), interacts with Tkv and thereby prevents binding and phosphorylation of Mad by Tkv (198).

The *in vivo* relevance of Smad6 and Smad7 is shown in several systems. Smad6 and Smad7 were initially identified as genes induced by shear stress in vascular endothelial cells (204). Smad7 inhibits activin-induced dorsal mesoderm formation and interferes with inhibition of activin-mediated growth arrest and apoptosis in B-cells (206). Smad6 and Smad7 antagonize BMP-induced mesoderm development, have neural-inducing potencies in *Xenopus*, and inhibit BMP-induced growth arrest and apoptosis in B cells (200, 205–207).

Alternative functions for Smad6 and Smad7 may exist as well. In *Xenopus*, XSmad6 was found to be partially or completely restricted to the nuclei in most cells (207). In mammalian cells, Smad7 has been located in the nucleus in the absence of ligand, but rapidly accumulated in the cytoplasm after TGF- $\beta$  stimulation (202). In accordance with its importance for inhibition of R-Smad signaling, the inhibitory Smad carboxyl-terminal tail also harbors domains required for transport across the nuclear membrane (202). A nuclear localization of inhibitory Smads might be required to allow phosphorylation of R-Smads after receptor activation; the mechanism whereby Smad7 is translocated to the plasma membrane after TGF- $\beta$  receptor activation is unknown. An interesting possibility, which remains to be elucidated, is that inhibitory Smads in addition function in transcriptional regulation. Differential compartmentalization of the inhibitory Smads, in combination with induction of their expression after signaling by TGF- $\beta$  superfamily members, and the fact that inhibitory Smads can selectively eliminate

particular R-Smad pathways provide a tightly controlled cell-autonomous regulatory mechanism.

#### Modulation of Smad signaling by MAP kinase pathways

The mitogen-activated protein kinase kinase kinase TAK1 (TGF- $\beta$  activated kinase 1) has been shown to be phosphorylated and activated upon TGF- $\beta$  or BMP-4 stimulation (208), and can mediate transcriptional activation of a luciferase reporter driven by a TGF- $\beta$ -inducible element of the PAI-1 promoter containing three AP-1 sites. This effect is strongly enhanced by the TAK1-activator TAB1 (209). The *in vivo* relevance of the cooperative functioning of TAK1 and TAB1 was demonstrated in *Xenopus* mesoderm development in which TAK1 and TAB1 promoted ventral mesoderm induction and perturbed neural differentiation, thereby substituting BMP signaling (210). The human X-chromosome-linked inhibitor of apoptosis protein (XIAP) can interact with both BMP receptors as well as with TAB1 and enhances ventralization of *Xenopus* embryos in a TAB1-TAK1-dependent manner (211).

Using the transmembrane and cytoplasmic domain of BMPR-IA as bait in yeast two-hybrid screens, BRAM1 (BMP receptor-associated molecule) was identified. The carboxyl-terminal region of BRAM1 is responsible for specific interaction with BMPR-IA and does not associate with the kinase domain of T $\beta$ R-I (212). Furthermore, this region in BRAM1 is also responsible for interaction with TAB1 (212). Expression of GST-coupled BMPR-IA, HA-tagged BRAM1, and myc-tagged TAB1 in COS7 cells showed that the three proteins form a ternary complex. BRAM1 might function as an adaptor protein for positioning TAB1 in close proximity of BMPR-IA kinase domain (212). The role of BRAM1 in activation of the TAB1/TAK1 signaling cascade, as well as its possible effect on Smad-receptor interaction and Smad activation is not clear at present.

A downstream phosphorylation target in the MAPK cascade triggered by TAK1 is mitogen-activated protein kinase kinase 4 (MKK4)/stress-activated protein kinase/extracellular signal-regulated kinase SEK1 (208, 213), which is involved in the stress-activated protein kinase (SAPK)/c-Jun NH<sub>2</sub>-terminal kinase (JNK) pathway, finally leading to activation of c-Jun. In that JNK activity induced by hematopoietic progenitor kinase-1 (HPK-1) can be perturbed by expression of a kinase inactive form of TAK1, there are indications that TGF- $\beta$  might exert JNK activation via the HPK1-TAK1-SAPK/JNK route (214). The small Rho-like GTPases Rho, Rac, and cdc42, which signal via the SAPK/JNK pathway, have also been shown to be important for certain aspects of TGF- $\beta$  signaling (215, 216).

The SAPK/JNK pathway has been shown to affect TGF- $\beta$  signaling via the Smad-pathway, since overexpression of dominant negative members of the SAPK/JNK pathway prevent both TGF- $\beta$ - and Smad4-mediated transcriptional response (217). The interface of SAPK/JNK- and Smad-signaling might be at the level of interaction between c-Jun and Smad3, since these transcription factors have been shown to associate after TGF- $\beta$  stimulation and synergize in activating a transcriptional reporter containing AP-1 binding sites (163, 166).

Other levels of interaction between MAPK pathways and Smad signaling have been elucidated. In the linker regions of R-Smads that connect the MH1 DNA binding domain with the MH2 transcriptional domain of receptor-activated Smads, several ERK recognition sites (PXSP, or PXTTP motifs) (Fig. 3B) and JNK motifs (XXSP) are present in the linker region of R-Smads. Phosphorylation of these sites in Smad1 by EGF or HGF has been shown to circumvent nuclear translocation of Smad1 and indicates a role for ERK signaling in the modulation of TGF- $\beta$  signaling (218). On the other hand, EGF and HGF have been shown to mediate signaling via Smad2, activated by kinases downstream of MEK1 (219). Furthermore, deletion of the carboxyl-terminal phosphorylation motif SS(V/M)S abrogates HGF-induced Smad1 or Smad2 phosphorylation, their nuclear translocation, and HGF-induced activation of a Smad responsive reporter construct, indicating that these residues are important for Smad functioning in both TGF- $\beta$  and HGF or EGF signaling (219). It is not known under which conditions the ERK signaling inhibits TGF- $\beta$  family/Smad-induced responses or mediates Smad-dependent pathways.

Analogously, differences in cellular context might underlie the apparently contradicting reports on the role of Ras/MAPK in TGF- $\beta$  signaling in different tumor cell systems. Phosphorylation of ERK sites in the linker region of Smad3 by activated Ras, MEK1, or v-HA-Ras-transformed mouse mammary epithelial Eph4 cells inhibits TGF- $\beta$ /Smad-mediated transcriptional responses, Smad nuclear translocation, and growth inhibition (220). In contrast, TGF- $\beta$  rapidly induces Ras/MEK signaling in intestinal epithelial IEC14 cells; this signal transduction pathway cooperates in TGF- $\beta$  mediated Smad-signaling, presumably by phosphorylation of Smad1 on the four ERK sites in the linker region (221).

## CONCLUSIONS AND PERSPECTIVES

Recent genetic and biochemical studies in *C. elegans*, *Drosophila*, *Xenopus*, and mammals have now firmly established the TGF- $\beta$ /Smad pathway as a pivotal means for intracellular signaling of TGF- $\beta$  superfamily

members (reviewed in refs 29, 101). On ligand-induced heteromeric complex formation of distinct type I and type II receptors, particular R-Smads transiently interact with and become phosphorylated by the activated type I receptor kinase. Subsequently, R-Smads assemble with Co-Smads into heteromeric complexes that translocate into the nucleus, where (in combination with other transcription factors) they regulate the transcription of target genes.

One TGF- $\beta$  superfamily member can couple with multiple type I receptors and multiple Smads, each mediating a distinct set of responses (reviewed in refs 29, 101). Thus, the receptor and Smad expression profile in the target cell is an important factor that decides which particular cellular responses are induced by a TGF- $\beta$  superfamily member. In addition, the repertoire of transcription factors that are present in the cell with which the Smads can interact is a critical determinant. Although multimerized Smad binding element is sufficient to drive TGF- $\beta$ -induced transcription (reviewed in ref 155), emerging promoter analyses of TGF- $\beta$  superfamily target genes indicate that transcriptional regulation is achieved by cooperation of Smads with other transcription factors. Complex formation of Smads with these transcription factors may occur independent of DNA binding or require DNA binding for additional selectivity (reviewed in ref 155).

The TGF- $\beta$ /Smad pathway has been implicated in many responses, including growth inhibition, differentiation and many transcriptional responses. An important issue that needs to be addressed is the requirement of particular Smads in these responses. In addition, it will be important to explore the possibility that Smad-independent signaling also occurs. TGF- $\beta$ -induced transcription of the fibronectin gene can occur independent of Smad4, but requires the JNK pathway and activation of cAMP-responsive element by c-Jun/ATF2 heterodimer (222).

An important task for the future is to determine the genetic programs that are triggered on challenging different cell types with different TGF- $\beta$  family members. This will become feasible by applying the newly developed functional genomics and proteomics technology and thereby determine the time- and dose-dependent effects of TGF- $\beta$  family members on the expression of thousands of genes/proteins simultaneously. In addition, changes in the expression patterns on ectopic expression of constitutively active or dominant negative versions of signaling components or on ligand stimulation in cells deficient in such components can be measured. The target genes/proteins identified will need experimental follow-up to evaluate their importance in various biological responses. Moreover, the involvement of different type I receptors and Smad molecules in



various responses induced by TGF- $\beta$  family member can be elucidated using this methodology.

The multifunctional character of TGF- $\beta$  superfamily members implicates a need for tight control of their activities. Indeed, both positive and negative (feedback) regulatory mechanisms have been observed at nearly every step in the TGF- $\beta$  superfamily signaling cascade, from the release of biologically active ligand to the Smad-mediated transcriptional effects. Cross talk of the TGF- $\beta$ /Smad pathway with other pathways will be a theme of many future studies, e.g., how other signaling inputs affect the activity, expression, stability, or subcellular localization of TGF- $\beta$  superfamily receptors and Smads. In addition, the pleiotropic action of TGF- $\beta$  superfamily members has driven the evolution of multiple closely related TGF- $\beta$  superfamily members with different expression and activation patterns. The mammalian TGF- $\beta$  superfamily will likely continue to grow until the complete human sequence has been elucidated at the beginning of the next century.

Much of our insight into signaling mechanisms has come from the genetically accessible model organisms *Drosophila* and *C. elegans* (reviewed in refs 29, 101). In *Drosophila*, dpp signals through two type I receptors—saxophone and thick veins—and their downstream effector, Mad. Recently, an activin/TGF- $\beta$  signal transduction pathway was identified in *Drosophila* (223, 224). Activation of the activin type I receptor Atr-I stimulates dSmad2-dependent pathways. The type II receptor punt and Medea are shared components between the Dpp and activin/TGF- $\beta$  pathway. *C. elegans* research obtained a boost by the recent completion of its entire genome sequence (reviewed in ref 225). This vertebrate has four TGF- $\beta$ -like ligands (including Daf-7 and Daf-1), one type II receptor (Daf-4), and two type I receptors (Daf-1 and Sma-6). Thus, Daf-4 is required for both pathways and type I receptors to determine signaling specificity (226). Daf-1 is important in dauer larva formation and Sma-6 for body size determination and male tail development. Six Smad proteins are present in the *C. elegans* genome: Sma-2, Sma-3, and Sma-4 have been implicated in the Daf-7/Daf-1 pathway, and the Smad mediators Daf-8, Daf-14, and Daf-3 in Dbl-1/Sma-6-induced signaling responses. *Drosophila* and *C. elegans* will continue to be extremely important in elucidating the molecular mechanisms that underlie TGF- $\beta$  superfamily signaling.

In mammals, the physiological significance of interactions between ligands and receptors, between receptors and Smads, and between Smads and Smad-interacting proteins, as well as the importance of the signaling pathways in which they act, need to be validated through comparison of the phenotypes of mice deficient in a particular TGF- $\beta$  superfamily member, receptor, Smad, or target gene. In those

cases where a null mutation leads to an early embryonal lethal phenotype, conditional knock-out approaches will be required to study the role of this component in late development or adult tissues. In addition, transgenic approaches in mice with dominant negative or constitutively active forms of TGF- $\beta$  superfamily signal transducers under inducible or tissue-specific promoters will provide important information regarding issues of signaling specificity and diversity in different cellular contexts.

The improved understanding by which TGF- $\beta$  superfamily members elicit and regulate different responses will be essential in the design of new therapeutic approaches for various diseases caused by deregulated activity of TGF- $\beta$  family members. For example, antagonists of TGF- $\beta$  could be applied in various types of fibrosis that are due to TGF- $\beta$  overactivity and agonists of TGF- $\beta$  in diseases in which enhanced activity is beneficial, such as wound healing or immunosuppression. In most cases it will be advantageous not to inhibit or activate all activities of TGF- $\beta$  superfamily members. By understanding the molecular mechanisms that underlie specificity, diversity, and regulation in TGF- $\beta$  superfamily signaling, it will be possible to screen for pharmacological compounds that inhibit or mimic only defined activities of TGF- $\beta$  superfamily members. □

We are grateful to our colleagues who generously contributed data prior to publication. We apologize to those whose contributions have not been cited due to space constraints.

## REFERENCES

1. Roberts, A. B., and Sporn, M. B. (1990) The transforming growth factor- $\beta$ s. In: *Peptide Growth Factors and Their Receptors, Part 1*, (Sporn, M. B., and Roberts, A. B., eds) Vol. 95, pp. 419–472, Springer-Verlag, Berlin.
2. Letterio, J. J., and Roberts, A. B. (1998) Regulation of immune responses by TGF- $\beta$ . *Annu. Rev. Immunol.* 16, 137–161.
3. Kingsley, D. M. (1994) The TGF- $\beta$  superfamily: new members, new receptors, and new genetic tests of function in different organisms. *Genes Dev.* 8, 133–146.
4. Hogan, B. L. M. (1996) Bone morphogenetic proteins: multifunctional regulators of vertebrate development. *Genes Dev.* 10, 1580–1594.
5. Mathews, I. S. (1994) Activin receptors and cellular signaling by the receptor serine kinase family. *Endocr. Rev.* 15, 310–325.
6. Roberts, A. B., and Sporn, M. B. (1992) Differential expression of the TGF- $\beta$  isoforms in embryogenesis suggests specific roles in developing and adult tissues. *Mol. Reprod. Dev.* 32, 91–98.
7. Shull, M. M., Ormsby, I., Kier, A. B., Pawlowski, S., Diebold, R. J., Yin, M., Allen, R., Sidman, C., Proetzel, G., Calvin, D., Annunziata, N., and Doetschman, T. (1992) Targeted disruption of the mouse transforming growth factor- $\beta$ 1 gene results in multifocal inflammatory disease. *Nature (London)* 359, 693–699.
8. Kulkarni, A. B., Huh, C.-G., Becker, D., Geiser, A., Lyght, M., Flanders, K. C., Roberts, A. B., Sporn, M. B., Ward, J. M., and Karlsson, S. (1993) Transforming growth factor  $\beta$ 1 null mutation in mice causes excessive inflammatory response and early death. *Proc. Natl. Acad. Sci. USA* 90, 770–774.
9. Diebold, R. J., Eis, M. J., Yin, M., Ormsby, I., Boivin, G. P., Darrow, B. J., Saffitz, J. E., and Doetschman, T. (1995) Early-onset multifocal inflammation in the transforming growth

- factor  $\beta 1$  null mouse is lymphocyte mediated. *Proc. Natl. Acad. Sci. USA* **92**, 12215-12219.
10. Dickson, M. C., Martin, J. S., Cousins, F. M., Kulkarni, A. B., Karlsson, S., and Akhurst, R. J. (1995) Defective haematopoiesis and vasculogenesis in transforming growth factor- $\beta 1$  knock out mice. *Development* **121**, 1845-1854.
  11. Letterio, J. J., Geiser, A. G., Kulkarni, A. B., Roche, N. S., Sporn, M. B., and Roberts, A. B. (1994) Maternal rescue of transforming growth factor- $\beta 1$  null mice. *Science* **264**, 1936-1938.
  12. Sanford, L. P., Ormsby, I., Gittenberger-de Groot, A. C., Sariola, H., Friedman, R., Boivin, G. P., Cardell, E. L., and Doetschman, T. (1997) TGF $\beta 2$  knockout mice have multiple developmental defects that are non-overlapping with other TGF $\beta$  knockout phenotypes. *Development* **124**, 2659-2670.
  13. Kaartinen, V., Voncken, J. W., Shuler, C., Warburton, D., Bu, D., Heisterkamp, N., and Groffen, J. (1995) Abnormal lung development and cleft palate in mice lacking TGF- $\beta 3$  indicates defects of epithelial-mesenchymal interaction. *Nat. Genet.* **11**, 415-421.
  14. Proetzel, G., Pawlowski, S. A., Wiles, M. V., Yin, M. Y., Boivin, G. P., Howles, P. N., Ding, J. X., Ferguson, M. W. J., and Doetschman, T. (1995) Transforming growth factor- $\beta 3$  is required for secondary palate fusion. *Nat. Genet.* **11**, 409-414.
  15. Roberts, A. B., Kim, S. J., Noma, T., Glick, A. B., Lafyatis, R., Lechleider, R., Jakowlew, S. B., Geiser, A., O'Reilly, M. A., Danielpour, D., et al. (1991) Multiple forms of TGF- $\beta$ : distinct promoters and differential expression. *Ciba Found. Symp.* **157**, 7-28.
  16. Kim, S. J., Park, K., Koeller, D., Kim, K. Y., Wakefield, L. M., Sporn, M. B., and Roberts, A. B. (1992) Post-transcriptional regulation of the human transforming growth factor- $\beta 1$  gene. *J. Biol. Chem.* **267**, 13702-13707.
  17. Munger, J. S., Harpel, J. G., Gleizes, P. E., Mazzieri, R., Nunes, I., and Rifkin, D. B. (1997) Latent transforming growth factor- $\beta$ : structural features and mechanisms of activation. *Kidney Int.* **51**, 1376-1382.
  18. Cui, Y., Jean, F., Thomas, G., and Christian, J. L. (1998) BMP-4 is proteolytically activated by furin and/or PC6 during vertebrate embryonic development. *EMBO J.* **17**, 4735-4743.
  19. Taipale, J., and Keski-Oja, J. (1997) Growth factors in the extracellular matrix. *FASEB J.* **11**, 51-59.
  20. Saharinen, J., Taipale, J., Monni, O., and Keski-Oja, J. (1998) Identification and characterization of a new latent transforming growth factor- $\beta$ -binding protein, LTBP-4. *J. Biol. Chem.* **273**, 18459-18469.
  21. Munger, J. S., Harpel, J. G., Giancotti, F. G., and Rifkin, D. B. (1998) Interactions between growth factors and integrins: latent forms of transforming growth factor- $\beta$  are ligands for the integrin  $\alpha \beta 1$ . *Mol. Biol. Cell* **9**, 2627-2638.
  22. Angeles-Cano, E. (1997) Structural basis for the pathophysiology of lipoprotein(a) in the athero-thrombotic process. *Braz. J. Med. Biol. Res.* **30**, 1271-1280.
  23. Miyazono, K., and Heldin, C. H. (1989) Role for carbohydrate structures in TGF- $\beta 1$  latency. *Nature (London)* **338**, 158-160.
  24. Barcellos-Hoff, M. H., and Dix, T. A. (1996) Redox-mediated activation of latent transforming growth factor- $\beta 1$ . *Mol. Endocrinol.* **10**, 1077-1083.
  25. Jullien, P., Berg, T. M., and Lawrence, D. A. (1989) Acidic cellular environments: activation of latent TGF- $\beta$  and sensitization of cellular responses to TGF- $\beta$  and EGF. *Int. J. Cancer* **43**, 886-891.
  26. Crawford, S. E., Stellmach, V., Murphy-Ullrich, J. E., Ribeiro, S. M. F., Lawler, J., Hynes, R. O., Boivin, G. P., and Boucek, N. (1998) Thrombospondin-1 is a major activator of TGF- $\beta 1$  in vivo. *Cell* **93**, 1159-1170.
  27. Cheifetz, S., Weatherbee, J. A., Tsang, M. L.-S., Anderson, J. K., Mole, J. E., Lucas, R., and Massagué, J. (1987) The transforming growth factor- $\beta$  system, a complex pattern of cross-reactive ligands and receptors. *Cell* **48**, 409-415.
  28. Ogawa, Y., Schmidt, D. K., Dasch, J. R., Chang, R.-J., and Glaser, C. B. (1992) Purification and characterization of transforming growth factor- $\beta 2.3$  and - $\beta 1.2$  heterodimers from bovine bone. *J. Biol. Chem.* **267**, 2325-2328.
  29. Massagué, J. (1998) TGF- $\beta$  signal transduction. *Annu. Rev. Biochem.* **67**, 753-791.
  30. Nishimatsu, S., and Thomsen, G. H. (1998) Ventral mesoderm induction and patterning by bone morphogenetic protein heterodimers in *Xenopus* embryos. *Mech. Dev.* **74**, 75-88.
  31. Israel, D. I., Nove, J., Kerns, K. M., Kaufman, R. J., Rosen, V., Cox, K. A., and Wozney, J. M. (1996) Heterodimeric bone morphogenetic proteins show enhanced activity *in vitro* and *in vivo*. *Growth Factors* **13**, 291-300.
  32. Sampath, T. K., Coughlin, J. E., Whetstone, R. M., Banach, D., Corbett, C., Ridge, R. J., Ozkaynak, E., Oppermann, H., and Rueger, D. C. (1990) Bovine osteogenic protein is composed of dimers of OP-1 and BMP-2A, two members of the transforming growth factor- $\beta$  superfamily. *J. Biol. Chem.* **265**, 13198-13205.
  33. Yamaguchi, Y., Mann, D. M., and Ruoslahti, E. (1990) Negative regulation of transforming growth factor- $\beta$  by the proteoglycan decorin. *Nature (London)* **346**, 281-284.
  34. Xu, T., Bianco, P., Fisher, L. W., Longenecker, G., Smith, E., Goldstein, S., Bonadio, J., Boskey, A., Heegaard, A. M., Sommer, B., Satomura, K., Dominguez, P., Zhao, C., Kulkarni, A. B., Robey, P. G., and Young, M. F. (1998) Targeted disruption of the biglycan gene leads to an osteoporosis-like phenotype in mice. *Nat. Genet.* **20**, 78-82.
  35. LaMarre, J., Hayes, M. A., Wollenberg, G. K., Hussaini, I., Hall, S. W., and Gonias, S. L. (1991) An  $\alpha_2$ -macroglobulin receptor-dependent mechanism for the plasma clearance of transforming growth factor- $\beta 1$  in mice. *J. Clin. Invest.* **87**, 39-44.
  36. Mather, J. P. (1996) Follistatins and  $\alpha 2$ -macroglobulin are soluble binding proteins for inhibin and activin. *Horm. Res.* **45**, 207-210.
  37. Fava, R. A., and McClure, D. B. (1987) Fibronectin-associated transforming growth factor. *J. Cell. Physiol.* **131**, 184-189.
  38. Paralkar, V. M., Vukicevic, S., and Reddi, A. H. (1991) Transforming growth factor  $\beta$  type 1 binds to collagen IV of basement membrane matrix: Implications for development. *Dev. Biol.* **143**, 303-308.
  39. François, V., and Bier, E. (1995) *Xenopus* chordin and *Drosophila* short gastrulation genes encode homologous proteins functioning in dorsal-ventral axis formation. *Cell* **80**, 19-20.
  40. Thomsen, G. H. (1997) Antagonism within and around the organizer: BMP inhibitors in vertebrate body patterning. *Trends Genet.* **13**, 209-211.
  41. Brunet, L. J., McMahon, J. A., McMahon, A. P., and Harland, R. M. (1998) Noggin, cartilage morphogenesis, and joint formation in the mammalian skeleton. *Science* **280**, 1455-1457.
  42. Hsu, D. R., Economides, A. N., Wang, X., Eimon, P. M., and Harland, R. M. (1998) The *Xenopus* dorsalizing factor gremlin identifies a novel family of secreted proteins that antagonize BMP activities. *Mol. Cell* **1**, 673-683.
  43. Iemura, S. I., Yamamoto, T. S., Takagi, C., Uchivama, H., Natsume, T., Shimazaki, S., Sugino, H., and Ueno, N. (1998) Direct binding of follistatin to a complex of bone morphogenetic protein and its receptor inhibits ventral and epidermal cell fates in early *Xenopus* embryo. *Proc. Natl. Acad. Sci. USA* **95**, 9337-9342.
  44. Blader, P., Rastegar, S., Fischer, N., and Strähle, U. (1997) Cleavage of the BMP-4 antagonist chordin by zebrafish tollid. *Science* **278**, 1937-1940.
  45. Mullins, M. C. (1998) Holy Tollid: tollid cleaves SOG/chordin to free DPP/BMPs. *Trends Genet.* **14**, 127-129.
  46. López-Casillas, F., Cheifetz, S., Doody, J., Andrés, J. L., Lane, W. S., and Massagué, J. (1991) Structure and expression of the membrane proteoglycan betaglycan, a component of the TGF- $\beta$  receptor system. *Cell* **67**, 785-795.
  47. Cheifetz, S., Bellón, T., Calés, C., Vera, S., Bernabeu, C., Massagué, J., and Letarte, M. (1992) Endoglin is a component of the transforming growth factor- $\beta$  receptor system in human endothelial cells. *J. Biol. Chem.* **267**, 19027-19030.
  48. Kaname, S., and Ruoslahti, E. (1996) Betaglycan has multiple binding sites for transforming growth factor- $\beta 1$ . *Biochem. J.* **315**, 815-820.
  49. López-Casillas, F., Wrana, J. L., and Massagué, J. (1993) Betaglycan presents ligand to the TGF $\beta$  signaling receptor. *Cell* **73**, 1435-1444.
  50. Sankar, S., Mahooti-Brooks, N., Centrella, M., McCarthy, T. L., and Madri, J. A. (1995) Expression of transforming growth factor  $\beta$  type III receptor in vascular endothelial cells increases their responsiveness to transforming growth factor  $\beta 2$ . *J. Biol. Chem.* **270**, 13567-13572.

51. López-Casillas, F., Payne, H. M., Andres, J. L., and Massagué, J. (1994) Betaglycan can act as a dual modulator of TGF- $\beta$  access to signaling receptors: Mapping of ligand binding and GAG attachment sites. *J. Cell Biol.* **124**, 557-568.
52. Barbara, N. P., Wrana, J. L., and Letarte, M. (1999) Endoglin is an accessory protein that interacts with the signaling receptor complex of multiple members of the transforming growth factor- $\beta$  superfamily. *J. Biol. Chem.* **274**, 584-594.
53. Lastres, P., Letamendia, A., Zhang, H. W., Rius, C., Almqvist, N., Raab, U., López, L. A., Langa, C., Fabra, A., Letarte, M., and Bernabéu, C. (1996) Endoglin modulates cellular responses to TGF- $\beta$ 1. *J. Cell Biol.* **133**, 1109-1121.
54. McAllister, K. A., Grogg, K. M., Johnson, D. W., Gallione, C. J., Baldwin, M. A., Jackson, C. E., Helmbold, E. A., Markel, D. S., McKinnon, W. C., Murrell, J., McCormick, M. K., Pericak-Vance, M. A., Heutink, P., Oostra, B. A., Haitjema, T., Westerman, C. J. J., Porteous, M. E., Guttacher, A. E., Letarte, M., and Marchuk, D. A. (1994) Endoglin, a TGF- $\beta$  binding protein of endothelial cells, is the gene for hereditary haemorrhagic telangiectasia type 1. *Nat. Genet.* **8**, 345-351.
55. Johnson, D. W., Berg, J. N., Baldwin, M. A., Gallione, C. J., Marondel, I., Yoon, S.-J., Stenzel, T. T., Speer, M., Pericak-Vance, M. A., Diamond, A., Guttacher, A. E., Jackson, C. E., Attisano, L., Kucherlapati, R., Porteous, M. E. M., and Marchuk, D. A. (1996) Mutations in the activin receptor-like kinase 1 gene in hereditary haemorrhagic telangiectasia type 2. *Nat. Genet.* **13**, 189-195.
56. Ohta, M., Greenberger, J. S., Anklesaria, P., Bassols, A., and Massagué, J. (1987) Two forms of transforming growth factor- $\beta$  distinguished by multipotential haematopoietic progenitor cells. *Nature (London)* **329**, 539-541.
57. Flanders, K. C., Lüdecke, C., Engels, S., Gissel, D. S., Roberts, A. B., Kondaiah, P., Lafyatis, R., Sporn, M. B., and Unsicker, K. (1991) Localization and actions of transforming growth factor- $\beta$ s in the embryonic nervous system. *Development* **113**, 183-191.
58. Shah, M., Foreman, D. M., and Ferguson, M. W. J. (1995) Neutralisation of TGF- $\beta$ 1 and TGF- $\beta$ 2 or exogenous addition of TGF- $\beta$ 3 to cutaneous rat wounds reduces scarring. *J. Cell. Sci.* **108**, 985-1002.
59. Cheifetz, S., and Massagué, J. (1991) Isoform-specific transforming growth factor- $\beta$  binding proteins with membrane attachments sensitive to phosphatidylinositol-specific phospholipase C. *J. Biol. Chem.* **266**, 20767-20772.
60. Mackay, K., and Danielpour, D. (1991) Novel 150- and 180-kDa glycoproteins that bind transforming growth factor (TGF)- $\beta$ 1 but not TGF- $\beta$ 2 are present in several cell lines. *J. Biol. Chem.* **266**, 9907-9911.
61. Bützow, R., Fukushima, D., Twardzik, D. R., and Ruoslahti, E. (1993) A 60-kD protein mediates the binding of transforming growth factor- $\beta$  to cell surface and extracellular matrix proteoglycans. *J. Cell Biol.* **122**, 721-727.
62. Pick, E., Franzén, P., Heldin, C.-H., and ten Dijke, P. (1997) Characterization of a 60-kDa cell surface-associated transforming growth factor- $\beta$  binding protein that can interfere with transforming growth factor- $\beta$  receptor binding. *J. Cell. Physiol.* **173**, 447-459.
63. Chen, R. H., Moses, H. L., Maruoka, E. M., Derynck, R., and Kawabata, M. (1995) Phosphorylation-dependent interaction of the cytoplasmic domains of the type I and type II transforming growth factor- $\beta$  receptors. *J. Biol. Chem.* **270**, 12235-12241.
64. Rodríguez, C., Chen, F., Weinberg, R. A., and Lodish, H. F. (1995) Cooperative binding of transforming growth factor (TGF)- $\beta$ 2 to the type I and II TGF- $\beta$  receptors. *J. Biol. Chem.* **270**, 15919-15922.
65. ten Dijke, P., Yamashita, H., Sampath, T. K., Reddi, A. H., Estevez, M., Riddle, D. L., Ichijo, H., Heldin, C.-H., and Miyazono, K. (1994) Identification of type I receptors for osteogenic protein-1 and bone morphogenetic protein-1. *J. Biol. Chem.* **269**, 16985-16988.
66. Rosenzweig, B. L., Imamura, T., Okadome, T., Cox, G. N., Yamashita, H., ten Dijke, P., Heldin, C.-H., and Miyazono, K. (1995) Cloning and characterization of a human type II receptor for bone morphogenetic proteins. *Proc. Natl. Acad. Sci. USA* **92**, 7632-7636.
67. Yamashita, H., ten Dijke, P., Franzén, P., Miyazono, K., and Heldin, C.-H. (1994) Formation of hetero-oligomeric complexes of type I and type II receptors for transforming growth factor- $\beta$ . *J. Biol. Chem.* **269**, 20172-20178.
68. Attisano, L., Cárcamo, J., Ventura, F., Weis, F., Massagué, J., and Wrana, J. L. (1993) Identification of human activin and TGF- $\beta$  type I receptors that form heteromeric kinase complexes with type II receptors. *Cell* **75**, 671-680.
69. Franzén, P., ten Dijke, P., Ichijo, H., Yamashita, H., Schulz, P., Heldin, C.-H., and Miyazono, K. (1993) Cloning of a TGF- $\beta$  type I receptor that forms a heteromeric complex with the TGF- $\beta$  type II receptor. *Cell* **75**, 681-692.
70. Attisano, L., Wrana, J. L., Cheifetz, S., and Massagué, J. (1992) Novel activin receptors: distinct genes and alternative mRNA splicing generate a repertoire of serine/threonine kinase receptors. *Cell* **68**, 97-108.
71. ten Dijke, P., Yamashita, H., Ichijo, H., Franzén, P., Laiho, M., Miyazono, K., and Heldin, C.-H. (1994) Characterization of type I receptors for transforming growth factor- $\beta$  and activin. *Science* **264**, 101-104.
72. Yamashita, H., ten Dijke, P., Huylebroeck, D., Sampath, T. K., Andries, M., Smith, J. C., Heldin, C.-H., and Miyazono, K. (1995) Osteogenic protein-1 binds to activin type II receptors and induces certain activin-like effects. *J. Cell Biol.* **130**, 217-226.
73. di Clemente, N., Wilson, C., Faure, E., Boussin, L., Camillo, P., Tizard, R., Picard, J.-Y., Vigier, B., Josso, N., and Cate, R. (1994) Cloning, expression, and alternative splicing of the receptor for anti-Müllerian hormone. *Mol. Endocrinol.* **8**, 1006-1020.
74. Rydén, M., Imamura, T., Jörnvall, H., Belluardo, N., Neveu, I., Trupp, M., Okadome, T., ten Dijke, P., and Ibáñez, C. F. (1996) A novel type I receptor serine-threonine kinase predominantly expressed in the adult central nervous system. *J. Biol. Chem.* **271**, 30603-30609.
75. Yelick, P. C., Abdulljabbar, T. S., and Stashenko, P. (1998) ALK-8, a novel type I serine/threonine kinase receptor, is expressed throughout early zebrafish development. *Dev. Dyn.* **211**, 352-361.
76. Renucci, A., Lemarchandel, V., and Rosa, F. (1996) An activated form of type I serine/threonine kinase receptor TARAM-A reveals a specific signalling pathway involved in fish head organizer formation. *Development* **122**, 3735-3743.
77. Wrana, J. L., Attisano, L., Wieser, R., Ventura, F., and Massagué, J. (1994) Mechanism of activation of the TGF- $\beta$  receptor. *Nature (London)* **370**, 341-347.
78. Feng, X.-H., and Derynck, R. (1997) A kinase subdomain of transforming growth factor- $\beta$  (TGF- $\beta$ ) type I receptor determines the TGF- $\beta$  intracellular signaling specificity. *EMBO J.* **16**, 3912-3923.
79. Chen, Y.-G., Hata, A., Lo, R. S., Wotton, D., Shi, Y.-G., Pavletich, N., and Massagué, J. (1998) Determinants of specificity in TGF- $\beta$  signal transduction. *Genes Dev.* **12**, 2144-2152.
80. Persson, U., Izumi, H., Soucheletsky, S., Itoh, S., Grimsby, S., Engström, U., Heldin, C.-H., Funa, K., and ten Dijke, P. (1998) The L45 loop in type I receptors for TGF- $\beta$  family members is a critical determinant in specifying Smad isoform activation. *FEBS Lett.* **434**, 83-87.
81. Luo, K.-X., and Lodish, H. F. (1997) Positive and negative regulation of type II TGF- $\beta$  receptor signal transduction by autophosphorylation on multiple serine residues. *EMBO J.* **16**, 1970-1981.
82. Koli, K. M., and Arteaga, C. L. (1997) Processing of the transforming growth factor- $\beta$  type I and II receptors: biosynthesis and ligand-induced regulation. *J. Biol. Chem.* **272**, 6423-6427.
83. Wells, R. G., Yankley, H., Lin, H. Y., and Lodish, H. F. (1997) Biosynthesis of the type I and type II TGF- $\beta$  receptors—implications for complex formation. *J. Biol. Chem.* **272**, 11444-11451.
84. Anders, R. A., Arline, S. L., Dor, J. J. E., and Leof, E. B. (1997) Distinct endocytic responses of heteromeric and homomeric transforming growth factor- $\beta$  receptors. *Mol. Biol. Cell* **8**, 2133-2143.
85. Anders, R. A., Dor, J. J. E., Arline, S. L., Garamszegi, N., and Leof, E. B. (1998) Differential requirement for type I and type II transforming growth factor- $\beta$  receptor kinase activity in ligand-mediated receptor endocytosis. *J. Biol. Chem.* **273**, 23118-23125.
86. Shoji, H., Nakamura, T., van den Eijnden-van Raaij, A. J. M., and Sugino, H. (1998) Identification of a novel type II activin



- receptor, type IIA-N, induced during the neural differentiation of murine P19 embryonal carcinoma cells. *Biochem. Biophys. Res. Commun.* **246**, 320-324
87. Xu, J., Matsuzaki, K., McKeen, K., Wang, F., Kan, M., and McKeen, W. L. (1994) Genomic structure and cloned cDNAs predict that four variants in the kinase domain of serine/threonine kinase receptors arise by alternative splicing and poly(A) addition. *Proc. Natl. Acad. Sci. USA* **91**, 7957-7961
  88. Suzuki, A., Shioda, N., Maeda, T., Tada, M., and Ueno, N. (1994) Cloning of an isoform of mouse TGF- $\beta$  type II receptor gene. *FEBS Lett.* **355**, 19-22
  89. Piek, E., Afrakhte, M., Sampath, K., van Zoelen, E. J., Heldin, C.-H., and ten Dijke, P. (1999) Functional antagonism between activin and osteogenic protein-1 in human embryonal carcinoma cells. *J. Cell. Physiol.* **180**, 141-149
  90. Lebrun, J. J., and Vale, W. W. (1997) Activin and inhibin have antagonistic effects on ligand-dependent heteromerization of the type I and type II activin receptors and human erythroid differentiation. *Mol. Cell. Biol.* **17**, 1682-1691
  91. Thisse, C., and Thisse, B. (1999) Antivin, a novel and divergent member of the TGF $\beta$  superfamily, negatively regulates mesoderm induction. *Development* **126**, 229-240
  92. Chen, R.-H., Miettinen, P. J., Maruoka, E. M., Choy, L., and Derynck, R. (1995) A WD-domain protein that is associated with and phosphorylated by the type II TGF- $\beta$  receptor. *Nature (London)* **377**, 548-552
  93. Choy, L., and Derynck, R. (1998) The type II transforming growth factor (TGF- $\beta$ ) receptor-interacting protein TRIP-1 acts as a modulator of the TGF- $\beta$  response. *J. Biol. Chem.* **273**, 31455-31462
  94. Datta, P. K., Chytil, A., Gorska, A. E., and Moses, H. L. (1998) Identification of STRAP, a novel WD domain protein in transforming growth factor- $\beta$  signaling. *J. Biol. Chem.* **273**, 34671-34674
  95. Griswold-Prenner, I., Kamibayashi, C., Maruoka, E. M., Mumby, M. C., and Derynck, R. (1998) Physical and functional interactions between type I transforming growth factor- $\beta$  receptors and Bx, a WD-40 repeat subunit of phosphatase 2A. *Mol. Cell. Biol.* **18**, 6595-6604
  96. Chang, M.-J., Zhang, D., Kinnunen, P., and Schneider, M. D. (1998) A novel protein distinguishes between quiescent and activated forms of the type I transforming growth factor- $\beta$  receptor. *J. Biol. Chem.* **273**, 9365-9368
  97. Wang, T., Li, B.-Y., Danielson, P. D., Shah, P. C., Rockwell, S., Lechleider, R. J., Martin, J., Manganaro, T., and Donahoe, P. K. (1996) The immunophilin FKBP12 functions as a common inhibitor of the TGF $\beta$  family type I receptors. *Cell* **86**, 435-444
  98. Chen, Y. G., Liu, F., and Massagué, J. (1997) Mechanism of TGF $\beta$  receptor inhibition by FKBP12. *EMBO J.* **16**, 3866-3876
  99. Wang, T., Danielson, P. D., Li, B.-Y., Shah, P. C., Kim, S. D., and Donahoe, P. K. (1996) The p21<sup>RAS</sup> farnesyltransferase  $\alpha$  subunit in TGF- $\beta$  and activin signaling. *Science* **271**, 1120-1122
  100. Ventura, F., Liu, F., Doody, J., and Massagué, J. (1996) Interaction of transforming growth factor- $\beta$  receptor I with farnesyl-protein transferase- $\alpha$  in yeast and mammalian cells. *J. Biol. Chem.* **271**, 13931-13934
  101. Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997) TGF- $\beta$  signalling from cell membrane to nucleus through SMAD proteins. *Nature (London)* **390**, 465-471
  102. Chen, D., Ji, X., Harris, M. A., Feng, J. Q., Karsenty, G., Celeste, A. J., Rosen, V., Mundy, G. R., and Harris, S. E. (1998) Differential roles for bone morphogenetic protein (BMP) receptor type IB and IA in differentiation and specification of mesenchymal precursor cells to osteoblast and adipocyte lineages. *J. Cell Biol.* **142**, 295-305
  103. Kawakami, Y., Ishikawa, T., Shimahara, M., Tanda, N., Enomoto-Iwamoto, M., Iwamoto, M., Kuwana, T., Ueki, A., Noji, S., and Nohno, T. (1996) BMP signaling during bone pattern determination in the developing limb. *Development* **122**, 3557-3566
  104. Zou, H., Wieser, R., Massagué, J., and Niswander, L. (1997) Distinct roles of type I bone morphogenetic protein receptors in the formation and differentiation of cartilage. *Genes Dev.* **11**, 2191-2203
  105. Nishitoh, H., Ichijo, H., Kimura, M., Matsumoto, T., Makishima, F., Yamaguchi, A., Yamashita, H., Enomoto, S., and Miyazono, K. (1996) Identification of type I and type II serine/threonine kinase receptors for growth/differentiation factor-5. *J. Biol. Chem.* **271**, 21345-21352
  106. Erlacher, L., McCartney, J., Piek, E., ten Dijke, P., Yanagishita, M., Oppermann, H., and Luyten, F. P. (1998) Cartilage-derived morphogenetic proteins and osteogenic protein-1 differentially regulate osteogenesis. *J. Bone Miner. Res.* **13**, 383-392
  107. Saitoh, M., Nishitoh, H., Amagasa, T., Miyazono, K., Takagi, M., and Ichijo, H. (1996) Identification of important regions in the cytoplasmic juxtamembrane domain of type I receptor that separate signaling pathways of transforming growth factor- $\beta$ . *J. Biol. Chem.* **271**, 2769-2775
  108. Souchevnytskyi, S., ten Dijke, P., Miyazono, K., and Heldin, C.-H. (1996) Phosphorylation of Ser165 in TGF- $\beta$  type I receptor modulates TGF- $\beta$ 1-induced cellular responses. *EMBO J.* **15**, 6231-6240
  109. Feng, X. H., Filvaroff, E. H., and Derynck, R. (1995) Transforming growth factor- $\beta$  (TGF- $\beta$ )-induced downregulation of cyclin A expression requires a functional TGF- $\beta$  receptor complex. Characterization of chimeric and truncated type I and type II receptors. *J. Biol. Chem.* **270**, 24237-24245
  110. Raftery, L. A., Twombly, V., Wharton, K., and Gelbart, W. M. (1995) Genetic screens to identify elements of the decapentaplegic signaling pathway in *Drosophila*. *Genetics* **139**, 241-254
  111. Sekelsky, J. J., Newfeld, S. J., Raftery, L. A., Chartoff, E. H., and Gelbart, W. M. (1995) Genetic characterization and cloning of *Moths against dpp*, a gene required for decapentaplegic function in *Drosophila melanogaster*. *Genetics* **139**, 1347-1358
  112. Savage, C., Das, P., Finelli, A. L., Townsend, S. R., Sun, C.-Y., Baird, S. E., and Padgett, R. W. (1996) *Caenorhabditis elegans* genes *smo-2*, *smo-3*, and *smo-4* define a conserved family of transforming growth factor- $\beta$  pathway components. *Proc. Natl. Acad. Sci. USA* **93**, 790-794
  113. Macías-Silva, M., Hoodless, P. A., Tang, S. J., Buchwald, M., and Wrana, J. L. (1998) Specific activation of Smad1 signaling pathways by the BMP7 type I receptor, ALK2. *J. Biol. Chem.* **273**, 25628-25636
  114. Chen, Y. G., and Massagué, J. (1999) Smad1 recognition and activation by the ALK1 group of transforming growth factor- $\beta$  family receptors. *J. Biol. Chem.* **274**, 3672-3677
  115. Thomsen, G. H. (1996) *Xenopus* *mothers against decapentaplegic* is an embryonic ventralizing agent that acts downstream of the BMP-2/4 receptor. *Development* **122**, 2359-2366
  116. Chen, Y., Bhushan, A., and Vale, W. (1997) Smad8 mediates the signaling of the receptor serine kinase. *Proc. Natl. Acad. Sci. USA* **94**, 12938-12943
  117. Tamaki, K., Souchevnytskyi, S., Itoh, S., Nakao, A., Sampath, K., Heldin, C.-H., and ten Dijke, P. (1998) Intracellular signaling of osteogenic protein-1 through Smad5 activation. *J. Cell Physiol.* **177**, 355-363
  118. Eppert, K., Scherer, S. W., Ozcelik, H., Pirone, R., Hoodless, P., Kim, H., Tsui, L.-C., Bapat, B., Gallinger, S., Andrulis, I. L., Thomsen, G. H., Wrana, J. L., and Attisano, L. (1996) MADR2 maps to 18q21 and encodes a TGF- $\beta$ -regulated MAD-related protein that is functionally mutated in colorectal carcinoma. *Cell* **86**, 543-552
  119. Macías-Silva, M., Abdollah, S., Hoodless, P. A., Pirone, R., Attisano, L., and Wrana, J. L. (1996) MADR2 is a substrate of the TGF $\beta$  receptor, and its phosphorylation is required for nuclear accumulation and signaling. *Cell* **87**, 1215-1224
  120. Zhang, Y., Feng, X.-H., Wu, R.-Y., and Derynck, R. (1996) Receptor-associated Mad homologues synergize as effectors of the TGF- $\beta$  response. *Nature (London)* **383**, 168-172
  121. Nakao, A., Imamura, T., Souchevnytskyi, S., Kawabata, M., Ishisaki, A., Oeda, E., Tamaki, K., Hanai, J.-I., Heldin, C.-H., Miyazono, K., and ten Dijke, P. (1997) TGF- $\beta$  receptor-mediated signalling through Smad2, Smad3, and Smad4. *EMBO J.* **16**, 5353-5362
  122. Baker, J. C., and Harland, R. M. (1996) A novel mesoderm inducer, Madr2, functions in the activin signal transduction pathway. *Genes Dev.* **10**, 1880-1889
  123. Graff, J. M., Bansal, A., and Melton, D. A. (1996) *Xenopus* Mad proteins transduce distinct subsets of signals for the TGF $\beta$  superfamily. *Cell* **85**, 479-487
  124. Nomura, M., and Li, E. (1998) Smad2 role in mesoderm formation, left-right patterning and craniofacial development. *Nature (London)* **393**, 786-790



125. Waldrip, W. R., Bikoff, E. K., Hoodless, P. A., Wrana, J. L., and Robertson, E. J. (1998) Smad2 signaling in extraembryonic tissues determines anterior-posterior polarity of the early mouse embryo. *Cell* **92**, 797-808.
126. Datto, M. B., Frederick, J. P., Pan, L., Borton, A. J., Zhuang, Y., and Wang, X.-F. (1999) Targeted disruption of Smad3 reveals an essential role in transforming growth factor- $\beta$  mediated signal transduction. *Mol. Cell. Biol.* **19**, 2495-2504.
127. Yang, X., Letterio, J. J., Lechleider, R. J., Chen, L., Hayman, R., Gu, H., Roberts, A. B., and Deng, C. (1999) Targeted disruption of Smad3 results in impaired mucosal immunity and diminished T cell responsiveness to TGF- $\beta$ . *EMBO J.* **18**, 1250-1291.
128. Zhu, Y., Richardson, J. A., Parada, L. F., and Graff, J. M. (1998) Smad3 mutant mice develop metastatic colorectal cancer. *Cell* **94**, 703-714.
129. Riggins, R. C., Kinzler, K. W., Vogelstein, B., and Thiagalingam, S. (1997) Frequency of Smad gene mutations in human cancers. *Cancer Res.* **57**, 2578-2580.
130. Liu, X.-J., Yue, J. B., Frey, R. S., Zhu, Q. C., and Mulder, K. M. (1998) Transforming growth factor  $\beta$  signaling through Smad1 in human breast cancer cells. *Cancer Res.* **58**, 4752-4757.
131. Bruno, E., Horrigan, S. K., Van Den Berg, D., Rozler, E., Fitting, P. R., Moss, S. T., Westbrook, C., and Hoffman, R. (1998) The Smad5 gene is involved in the intracellular signaling pathways that mediate the inhibitory effects of transforming growth factor- $\beta$  on human hematopoiesis. *Blood* **91**, 1917-1923.
132. Hahn, S. A., Schutte, M., Hoque, A. T. M. S., Moskaluk, C. A., da Costa, L. T., Rozenblum, E., Weinstein, C. L., Fischer, A., Yeo, C. J., Hruban, R. H., and Kern, S. E. (1996) DPC4, a candidate tumor suppressor gene at human chromosome 18q21.1. *Science* **271**, 350-353.
133. Das, P., Maduzia, L. L., Wang, H., Finelli, A. L., Cho, S. H., Smith, M. M., and Padgett, R. W. (1998) The *Drosophila* gene *Medea* demonstrates the requirement for different classes of Smads in dpp signaling. *Development* **125**, 1519-1528.
134. Lagna, G., Hata, A., Hemmati-Brivanlou, A., and Massagué, J. (1996) Partnership between DPC4 and SMAD proteins in TGF- $\beta$  signalling pathways. *Nature (London)* **383**, 832-836.
135. de Winter, J. P., Roelen, B. A. J., ten Dijke, P., van der Burg, B., and van den Eijnden-van Raaij, A. J. M. (1997) DPC4 (SMAD4) mediates transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1) induced growth inhibition and transcriptional response in breast tumour cells. *Oncogene* **14**, 1891-1899.
136. Liu, F., Pouppnot, C., and Massagué, J. (1997) Dual role of the Smad4/DPC4 tumor suppressor in TGF $\beta$ -inducible transcriptional complexes. *Genes Dev.* **11**, 3157-3167.
137. Zhang, Y., Musci, T., and Derynck, R. (1997) The tumor suppressor Smad4/DPC4 as a central mediator of Smad function. *Curr. Biol.* **7**, 270-276.
138. Schutte, M., Hruban, R. H., Hedrick, L., Cho, K. R., Nadasdy, C. M., Weinstein, C. L., Bova, G. S., Isaacs, W. B., Cairns, P., Nawroz, H., Sidransky, D., Casero, R. A., Jr., Meltzer, P. S., Hahn, S. A., and Kern, S. E. (1996) DPC4 gene in various tumor types. *Cancer Res.* **56**, 2527-2530.
139. Shi, Y., Hata, A., Lo, R. S., Massagué, J., and Pavletich, N. P. (1997) A structural basis for mutational inactivation of the tumour suppressor Smad4. *Nature (London)* **388**, 87-93.
140. Le Dai, J., Turnacioglu, K. K., Schutte, M., Sugar, A. Y., and Kern, S. E. (1998) Dpc4 transcriptional activation and dysfunction in cancer cells. *Cancer Res.* **58**, 4592-4597.
141. Zhou, S. B., Buckhaults, P., Zawel, L., Bunz, F., Riggins, G., Dai, J. L., Kern, S. E., Kinzler, K. W., and Vogelstein, B. (1998) Targeted deletion of Smad4 shows it is required for transforming growth factor  $\beta$  and activin signaling in colorectal cancer cells. *Proc. Natl. Acad. Sci. USA* **95**, 2412-2416.
142. Sirard, C., de la Pompa, J. L., Elia, A., Itie, A., Mirtsos, C., Cheung, A., Hahn, S., Wakeham, A., Schwartz, L., Kern, S. E., Rossant, J., and Mak, T. W. (1998) The tumor suppressor gene Smad4/Dpc4 is required for gastrulation and later for anterior development of the mouse embryo. *Genes Dev.* **12**, 107-119.
143. Wisotzkey, R. G., Mehra, A., Sutherland, D. J., Dobens, L. L., Liu, X., Dohrmann, C., Attisano, L., and Raftery, L. A. (1998) *Medea* is a *Drosophila* Smad 4 homolog that is differentially required to potentiate DPP responses. *Development* **125**, 1433-1445.
144. LeSueur, J. A., and Graff, J. M. (1999) Spemann organizer activity of Smad10. *Development* **126**, 137-146.
145. Masuyama, N., Hanafusa, H., Kusakabe, M., Shibuya, H., and Nishida, E. (1999) Identification of two Smad4 proteins in *Xenopus*. Their common and distinct properties. *J. Biol. Chem.* **274**, 12163-12170.
146. Kawabata, M., Inoue, H., Hanyu, A., Imamura, T., and Miyazono, K. (1998) Smad proteins exist as monomers *in vivo* and undergo homo- and hetero-oligomerization upon activation by serine/threonine kinase receptors. *EMBO J.* **17**, 4056-4065.
147. Tsukazaki, T., Chiang, T. A., Davison, A. F., Attisano, L., and Wrana, J. L. (1998) SARA, a FYVE domain protein that recruits Smad2 to the TGF $\beta$  receptor. *Cell* **95**, 779-791.
148. ten Dijke, P., and Heldin, C.-H. (1998) An anchor for activation. *Nature (London)* **397**, 109-111.
149. Lo, R. S., Chen, Y. G., Shi, Y., Pavletich, N. P., and Massagué, J. (1998) The L3 loop: a structural motif determining specific interactions between SMAD proteins and TGF- $\beta$  receptors. *EMBO J.* **17**, 996-1005.
150. Abdollah, S., Macías-Silva, M., Tsukazaki, T., Hayashi, H., Attisano, L., and Wrana, J. L. (1997) T $\beta$ RI phosphorylation of Smad2 on Ser<sup>165</sup> and Ser<sup>167</sup> is required for Smad2-Smad4 complex formation and signaling. *J. Biol. Chem.* **272**, 27678-27685.
151. Soucheletsky, S., Tamaki, K., Engström, U., Wernstedt, C., ten Dijke, P., and Heldin, C.-H. (1997) Phosphorylation of Ser<sup>165</sup> and Ser<sup>167</sup> in the C terminus of Smad2 mediates interaction with Smad4 and is required for transforming growth factor- $\beta$  signaling. *J. Biol. Chem.* **272**, 28107-28115.
152. Kretschmar, M., Liu, F., Hata, A., Doody, J., and Massagué, J. (1997) The TGF- $\beta$  family mediator Smad1 is phosphorylated directly and activated functionally by the BMP receptor kinase. *Genes Dev.* **11**, 984-995.
153. Wu, R.-Y., Zhang, Y., Feng, X.-H., and Derynck, R. (1997) Heteromeric and homomeric interactions correlate with signaling activity and functional cooperativity of Smad3 and Smad4/DPC4. *Mol. Cell. Biol.* **17**, 2521-2528.
154. Hata, A., Lo, R. S., Wotton, D., Lagna, G., and Massagué, J. (1997) Mutations increasing autoinhibition inactivate tumour suppressors Smad2 and Smad4. *Nature (London)* **388**, 82-87.
155. Derynck, R., Zhang, Y., and Feng, X.-H. (1998) Smads: Transcriptional activators of TGF- $\beta$  responses. *Cell* **95**, 737-740.
156. Kim, J., Johnson, K., Chen, H. J., Carroll, S., and Laughon, A. (1997) *Drosophila* Mad binds to DNA and directly mediates activation of *vestigial* by Decapentaplegic. *Nature (London)* **388**, 304-308.
157. Vindevoghel, L., Lechleider, R. J., Kon, A., de Caestecker, M. P., Uitto, J., Roberts, A. B., and Mauviel, A. (1998) SMAD3/4-dependent transcriptional activation of the human type VII collagen (COL7A1) promoter by transforming growth factor  $\beta$ . *Proc. Natl. Acad. Sci. USA* **95**, 14769-14774.
158. Yingling, J. M., Datto, M. B., Wong, C., Frederick, J. P., Liberati, N. T., and Wang, X.-F. (1997) Tumor suppressor Smad4 is a transforming growth factor  $\beta$ -inducible DNA binding protein. *Mol. Cell. Biol.* **17**, 7019-7028.
159. Jonk, L. J., Itoh, S., Heldin, C.-H., ten Dijke, P., and Kruijer, W. (1998) Identification and functional characterization of a Smad binding element (SBE) in the *JunB* promoter that acts as a transforming growth factor- $\beta$ , activin, and bone morphogenetic protein-inducible enhancer. *J. Biol. Chem.* **273**, 21145-21152.
160. Denlier, S., Itoh, S., Vivien, D., ten Dijke, P., Huet, S., and Gauthier, J.-M. (1998) Direct binding of Smad3 and Smad4 to critical TGF $\beta$ -inducible elements in the promoter of human plasminogen activator inhibitor-type 1 gene. *EMBO J.* **17**, 3091-3100.
161. Zawel, L., Dai, J. L., Buckhaults, P., Zhou, S., Kinzler, K. W., Vogelstein, B., and Kern, S. E. (1998) Human Smad3 and Smad4 are sequence-specific transcription activators. *Mol. Cell* **1**, 611-617.
162. Shi, Y., Wang, Y. F., Jayaraman, L., Yang, H., Massagué, J., and Pavletich, N. P. (1998) Crystal structure of a Smad MH1 domain bound to DNA: insights on DNA binding in TGF- $\beta$  signaling. *Cell* **94**, 585-594.
163. Yagi, K., Goto, D., Hamamoto, T., Takenoshita, S., Kato, M., and Miyazono, K. (1999) Alternatively-spliced variant of Smad2

- lacking exon 3: Comparison with wild-type Smad2 and Smad3. *J. Biol. Chem.* **274**, 703-709
164. Dennler, S., Huett, S., and Gauthier, J.-M. (1999) A short amino acid sequence in MH1 domain is responsible for functional differences between Smad2 and Smad3. *Oncogene* **18**, 1643-1648
165. Labbé, E., Silvestri, C., Hoodless, P. A., Wrana, J. L., and Attisano, L. (1998) Smad2 and Smad3 positively and negatively regulate TGF $\beta$ -dependent transcription through the forkhead DNA-binding protein FAST2. *Mol. Cell* **2**, 109-120
166. Zhang, Y., Feng, X. H., and Derynck, R. (1998) Smad3 and Smad4 cooperate with c-Jun/c-Fos to mediate TGF $\beta$ -induced transcription. *Nature (London)* **394**, 909-913
167. Zhou, S., Zawel, L., Lengauer, C., Kinzler, K. W., and Vogelstein, B. (1998) Characterization of human FAST-1, a TGF $\beta$  and activin signal transducer. *Mol. Cell* **2**, 121-127
168. Hua, X. X., Liu, X. D., Ansari, D. O., and Lodish, H. F. (1998) Synergistic cooperation of TFE3 and Smad proteins in TGF $\beta$ -induced transcription of the plasminogen activator inhibitor-1 gene. *Genes Dev* **12**, 3084-3095
169. Datto, M. B., Yu, Y., and Wang, X.-F. (1995) Functional analysis of the transforming growth factor  $\beta$  responsive elements in the WAF1/Cip1/p21 promoter. *J. Biol. Chem.* **270**, 28623-28628
170. Li, J.-M., Nichols, M. A., Chandrasekharan, S., Xiong, Y., and Wang, X.-F. (1995) Transforming growth factor  $\beta$  activates the promoter of cyclin-dependent kinase inhibitor p15<sup>INK4B</sup> through an Sp1 consensus site. *J. Biol. Chem.* **270**, 26750-26753
171. Moustakas, A., and Kardassis, D. (1998) Regulation of the human p21/WAF1/Cip1 promoter in hepatic cells by functional interactions between Sp1 and Smad family members. *Proc. Natl. Acad. Sci. USA* **95**, 6733-6738
172. Liberati, N. T., Datto, M. B., Frederick, J. P., Shen, X., Wong, C., Rougier-Chapman, E. M., and Wang, X.-F. (1999) Smads bind directly to the Jun family of AP-1 transcription factors. *Proc. Natl. Acad. Sci. USA* **96**, 4844-4849
173. Wong, C., Rougier-Chapman, E. M., Frederick, J. P., Datto, M. B., Liberati, N. T., Li, J.-M., and Wang, X.-F. (1999) Smad3/Smad4 and AP-1 complexes synergize in transcriptional activation of the c-Jun promoter by transforming growth factor- $\beta$ . *Mol. Cell Biol.* **19**, 1821-1830
174. Liu, B., Dou, C.-L., Prabhu, L., and Lai, E. (1999) FAST-2 is a mammalian winged-helix protein which mediates transforming growth factor  $\beta$  signals. *Mol. Cell Biol.* **19**, 424-430
175. Liu, F., Hata, A., Baker, J. C., Doody, J., Cárcamo, J., Harland, R. M., and Massagué, J. (1996) A human Mad protein acting as a BMP-regulated transcriptional activator. *Nature (London)* **381**, 620-623
176. Feng, X. H., Zhang, Y., Wu, R. Y., and Derynck, R. (1998) The tumor suppressor Smad4/DPC4 and transcriptional adaptor CBP/p300 are coactivators for Smad3 in TGF $\beta$ -induced transcriptional activation. *Genes Dev.* **12**, 2153-2163
177. Wrana, J., and Pawson, T. (1997) Mad about SMADs. *Nature (London)* **388**, 28-29
178. de Caestecker, M. P., Hemmati, P., Larisch-Bloch, S., Ajmura, R., Roberts, A. B., and Lechleider, R. J. (1997) Characterization of functional domains within Smad4/DPC4. *J. Biol. Chem.* **272**, 13690-13696
179. Shioda, T., Lechleider, R. J., Dunwoodie, S. L., Li, H., Yahata, T., de Caestecker, M. P., Fennier, M. H., Roberts, A. B., and Isselbacher, K. J. (1998) Transcriptional activating activity of Smad4: roles of SMAD hetero-oligomerization and enhancement by an associating transactivator. *Proc. Natl. Acad. Sci. USA* **95**, 9785-9790
180. Chen, X., Rubock, M. J., and Whitman, M. (1996) A transcriptional partner for MAD proteins in TGF $\beta$  signalling. *Nature (London)* **383**, 691-696
181. Chen, X., Weisberg, E., Fridmacher, V., Watanabe, M., Naco, G., and Whitman, M. (1997) Smad4 and FAST-1 in the assembly of activin-responsive factor. *Nature (London)* **389**, 85-89
182. Shimizu, A., Mitsuyasu, K., Nakao, A., Imamura, T., ten Dijke, P., Heldin, C.-H., Kawabata, M., Shimada, S., and Miyazono, K. (1998) Identification of receptors and Smad proteins involved in activin signalling in a human epidermal keratinocyte cell line. *Genes Cells* **3**, 125-134
183. Yanagisawa, K., Osada, H., Masuda, A., Kondo, M., Saito, T., Yatabe, Y., Takagi, K., Takahashi, T., and Takahashi, T. (1998) Induction of apoptosis by Smad3 and down-regulation of Smad3 expression in response to TGF $\beta$  in human small lung epithelial cells. *Oncogene* **17**, 1743-1747
184. Janknecht, R., Wells, N. J., and Hunter, T. (1998) TGF $\beta$ -stimulated cooperation of Smad proteins with the coactivators CBP/p300. *Genes Dev.* **12**, 2114-2119
185. Shen, X., Hu, P. P., Liberati, N. T., Datto, M. B., Frederick, J. P., and Wang, X. F. (1998) TGF $\beta$ -induced phosphorylation of Smad3 regulates its interaction with coactivator p300/CBP-binding protein. *Mol. Biol. Cell* **9**, 3309-3319
186. Topper, J. N., DiChiara, M. R., Brown, J. D., Williams, A. J., Falb, D., Collins, T., and Gimbrone, M. A., Jr. (1998) CREB binding protein is a required coactivator for Smad-dependent transforming growth factor  $\beta$  transcriptional responses in endothelial cells. *Proc. Natl. Acad. Sci. USA* **95**, 9506-9511
187. Nishihara, A., Hanai, J.-I., Okamoto, N., Yanagisawa, J., Kato, S., Miyazono, K., and Kawabata, M. (1998) Role of p300, a transcriptional coactivator, in signalling of TGF $\beta$ . *Genes Cells* **3**, 613-623
188. Abraham, S. E., Carter, M. C., and Moran, E. (1992) Transforming growth factor  $\beta$ 1 (TGF $\beta$ 1) reduces cellular levels of p34<sup>cdc2</sup>, and this effect is abrogated by adenovirus independently of the E1A-associated pRB binding activity. *Mol. Biol. Cell* **3**, 655-665
189. Datto, M. B., Hu, P. P. C., Kowalik, T. F., Yingling, J., and Wang, X. F. (1997) The viral oncoprotein E1A blocks transforming growth factor  $\beta$ -mediated induction of p21/WAF1/Cip1 and p15/INK4B. *Mol. Cell Biol.* **17**, 2030-2037
190. Candia, A. F., Watabe, T., Hawley, S. H. B., Onichtchouk, D., Zhang, Y., Derynck, R., Niehrs, C., and Cho, K.-W. Y. (1997) Cellular interpretation of multiple TGF $\beta$  signals: intracellular antagonism between activin/Bvg1 and BMP-2/4 signaling mediated by Smads. *Development* **124**, 4467-4480
191. Horvai, A. E., Xu, L., Korzus, E., Brard, G., Kalafus, D., Mullen, T. M., Rose, D. W., Rosenfeld, M. G., and Glass, C. K. (1997) Nuclear integration of JAK/STAT and Ras/AP-1 signaling by CBP and p300. *Proc. Natl. Acad. Sci. USA* **94**, 1074-1079
192. Kurokawa, M., Mitani, K., Irie, K., Matsuyama, T., Takahashi, T., Chiba, S., Yazaki, Y., Matsumoto, K., and Hirai, H. (1998) The oncoprotein Evi-1 represses TGF $\beta$  signalling by inhibiting Smad3. *Nature (London)* **394**, 92-96
193. Perkins, A. S., Mercer, J. A., Jenkins, N. A., and Copeland, N. G. (1991) Patterns of *Evi-1* expression in embryonic and adult tissues suggest that *Evi-1* plays an important regulatory role in mouse development. *Development* **111**, 479-487
194. Hayashi, H., Abdollahi, S., Qiu, Y., Cai, J., Xu, Y. Y., Grinnell, B. W., Richardson, M. A., Topper, J. N., Gimbrone, M. A. J., Wrana, J. L., and Falb, D. (1997) The MAD-related protein Smad7 associates with the TGF $\beta$  receptor and functions as an antagonist of TGF $\beta$  signaling. *Cell* **89**, 1165-1173
195. Imamura, T., Takase, M., Nishihara, A., Oeda, E., Hanai, J., Kawabata, M., and Miyazono, K. (1997) Smad6 inhibits signaling by the TGF $\beta$  superfamily. *Nature (London)* **389**, 622-626
196. Nakao, A., Afrakhte, M., Morén, A., Nakayama, T., Christian, J. L., Heuchel, R., Itoh, S., Kawabata, M., Heldin, N.-E., Heldin, C.-H., and ten Dijke, P. (1997) Identification of Smad7, a TGF $\beta$ -inducible antagonist of TGF $\beta$  signalling. *Nature (London)* **389**, 631-635
197. Tsuneizumi, K., Nakayama, T., Kamoshida, Y., Kornberg, T. B., Christian, J. L., and Tabata, T. (1997) Daughters against dpp modulates dpp organizing activity in Drosophila wing development. *Nature (London)* **389**, 627-631
198. Inoue, H., Imamura, T., Ishidou, Y., Takase, M., Udagawa, Y., Oka, Y., Tsuneizumi, K., Tabata, T., Miyazono, K., and Kawabata, M. (1998) Interplay of signal mediators of decapentaplegic (Dpp): molecular characterization of mothers against dpp, Medea, and daughters against dpp. *Mol. Biol. Cell* **9**, 2145-2156
199. Souchelnytskyi, S., Nakayama, T., Nakao, A., Morén, A., Heldin, C.-H., Christian, J. L., and ten Dijke, P. (1998) Physical and functional interaction of murine and *Xenopus* Smad7 with bone morphogenetic protein receptors and transforming growth factor- $\beta$  receptors. *J. Biol. Chem.* **273**, 25364-25370
200. Hata, A., Lagna, G., Massagué, J., and Hemmati-Brivanlou, A. (1998) Smad6 inhibits BMP/Smad1 signaling by specifically competing with the Smad4 tumor suppressor. *Genes Dev.* **12**, 186-197
201. Afrakhte, M., Morén, A., Jossan, S., Itoh, S., Sampath, K., Westermarck, B., Heldin, C.-H., Heldin, N.-E., and ten Dijke, P.



- (1998) Induction of inhibitory Smad6 and Smad7 mRNA by TGF- $\beta$  family members. *Biochem. Biophys. Res. Commun.* **249**, 505-511.
202. Itoh, S., Landstrom, M., Hermansson, A., Itoh, F., Heldin, C.-H., Heldin, N.-E., and ten Dijke, P. (1998) Transforming growth factor- $\beta$  induces nuclear export of inhibitory Smad7. *J. Biol. Chem.* **273**, 29195-29201.
203. Takase, M., Imamura, T., Sampath, T. K., Takeda, K., Ichijo, H., Miyazono, K., and Kawabata, M. (1998) Induction of Smad6 mRNA by bone morphogenetic proteins. *Biochem. Biophys. Res. Commun.* **244**, 26-29.
204. Topper, J. N., Cai, J., Qiu, Y., Anderson, K. R., Xu, Y. Y., Deeds, J. D., Feeley, T., Gimeno, C. J., Woolf, E. A., Tayber, O., Mays, G. C., Sampson, B. A., Schoen, F. J., Gimbrone, M. A., Jr., and Falb, D. (1997) Vascular *MADs*: two novel *MAD*-related genes selectively inducible by flow in human vascular endothelium. *Proc. Natl. Acad. Sci. USA* **94**, 9314-9319.
205. Nakayama, T., Snyder, M., Grewal, S., Tsuneizumi, K., Tabata, T., and Christian, J. L. (1998) Xenopus Smad8 acts downstream of RMP-4 to modulate its activity during vertebrate embryonic patterning. *Development* **125**, 857-867.
206. Ishiaki, A., Yamato, K., Hashimoto, S., Nakao, A., Tamaki, K., Nonaka, K., ten Dijke, P., Sugino, H., and Nishihara, T. (1998) Differential inhibition of Smad6 and Smad7 on bone morphogenetic protein- and activin-mediated growth arrest and apoptosis in B cells. *J. Biol. Chem.* **273**, 13637-13642.
207. Nakayama, T., Garner, H., Berg, L. K., and Christian, J. L. (1998) Smad6 functions as an intracellular antagonist of some TGF- $\beta$  family members during Xenopus embryogenesis. *Genes Cells* **3**, 387-394.
208. Yamaguchi, K., Shirakabe, K., Shibuya, H., Irie, K., Oishi, I., Ueno, N., Taniguchi, T., Nishida, E., and Matsumoto, K. (1995) Identification of a member of the MAPKKK family as a potential mediator of TGF- $\beta$  signal transduction. *Science* **270**, 2008-2011.
209. Shibuya, H., Yamaguchi, K., Shirakabe, K., Tonegawa, A., Gotoh, Y., Ueno, N., Irie, K., Nishida, E., and Matsumoto, K. (1996) TAB1, an activator of the TAK1/MAPKKK in TGF- $\beta$  signal transduction. *Science* **272**, 1179-1182.
210. Shibuya, H., Iwata, H., Masuyama, N., Gotoh, Y., Yamaguchi, K., Irie, K., Matsumoto, K., Nishida, E., and Ueno, N. (1998) Role of TAK1 and TAB1 in BMP signaling in early Xenopus development. *EMBO J.* **17**, 1019-1028.
211. Yamaguchi, K., Nagai, S.-I., Ninomiya-Tsuji, J., Nishida, E., Tamai, K., Irie, K., Ueno, N., Nishida, E., Shibuya, H., and Matsumoto, K. (1999) XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAK1 in the BMP signaling pathway. *EMBO J.* **19**, 1821-1830.
212. Kurozumi, K., Nishida, E., Yamaguchi, K., Fujita, T., Ueno, N., and Shibuya, H. (1998) BRAM1, a BMP receptor-associated molecule involved in BMP signalling. *Genes Cells* **3**, 257-264.
213. Shirakabe, K., Yamaguchi, K., Shibuya, H., Irie, K., Matsuda, S., Moriyuchi, T., Gotoh, Y., Matsumoto, K., and Nishida, E. (1997) TAK1 mediates the ceramide signaling to stress-activated protein kinase/c-Jun N-terminal kinase. *J. Biol. Chem.* **272**, 8141-8144.
214. Wang, W., Zhou, G., Hu, M.-C., Tsao, Z., and Jan, T.-H. (1997) Activation of the hematopoietic progenitor kinase-1 (HPK1)-dependent stress-activated c-Jun N-terminal kinase (JNK) pathway by transforming growth factor- $\beta$  (TGF- $\beta$ )-activated kinase (TAK1), a kinase mediator of TGF- $\beta$  signal transduction. *J. Biol. Chem.* **272**, 22771-22775.
215. Mucci, L., Skorecki, K. L., and Goldberg, H. J. (1996) Extracellular signal-regulated kinase and the small GTP-binding protein, Rac, contribute to the effects of transforming growth factor- $\beta$ 1 on gene expression. *J. Biol. Chem.* **271**, 16567-16572.
216. Atfi, A., Djelloul, S., Chastre, E., Davis, R. R., and Gaspach, C. (1997) Evidence for a role of Rho-like GTPases and stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) in transforming growth factor- $\beta$ -mediated signaling. *J. Biol. Chem.* **272**, 1429-1432.
217. Atfi, A., Buisine, M., Mazars, A., and Gaspach, C. (1997) Induction of apoptosis by DPC4, a transcriptional factor regulated by transforming growth factor- $\beta$  through stress-activated protein kinase/c-Jun N-terminal kinase (SAPK/JNK) signaling pathway. *J. Biol. Chem.* **272**, 24731-24734.
218. Kretschmar, M., Doody, J., and Massagué, J. (1997) Opposing BMP and EGF signalling pathways converge on the TGF- $\beta$  family mediator Smad1. *Nature (London)* **389**, 618-622.
219. De Caestecker, M. P., Parks, W. T., Frank, C. J., Castagnino, P., Bottaro, D. P., Roberts, A. B., and Lechleider, R. J. (1998) Smad2 transduces common signals from receptor serine-threonine and tyrosine kinases. *Genes Dev.* **12**, 1587-1592.
220. Kretschmar, M., Doody, J., Timokhina, L., and Massagué, J. (1999) A mechanism of repression of TGF- $\beta$ /Smad signaling by oncogenic Ras. *Genes Dev.* **13**, 804-816.
221. Yue, J., Frey, R. S., and Mulder, K. M. (1999) Cross-talk between the Smad1 and Ras/MEK signaling pathways for TGF- $\beta$ . *Oncogene* **58**, 4752-4757.
222. Hocoar, B. A., Brown, T. L., and Howe, P. H. (1999) TGF- $\beta$  induces fibronectin synthesis through a c-Jun N-terminal kinase-dependent, Smad4-independent pathway. *EMBO J.* **18**, 1345-1356.
223. Brummel, T., Abdollah, S., Haerry, T. E., Shimell, M. J., Merriam, J., Raftery, L., Wrana, J. L., and O'Connor, M. B. (1999) The Drosophila activin receptor Baboon signals through dSmad2 and controls cell proliferation but not patterning during larval development. *Genes Dev.* **13**, 98-111.
224. Das, P., Inoue, H., Baker, J. C., Beppu, H., Kawabata, M., Harland, R. M., Miyazono, K., and Padgett, R. W. (1999) Drosophila DSmad2 and Atr-1 transmit activin/TGF- $\beta$  signals. *Genes Cells* **4**, 123-134.
225. Ruvkun, G., and Hobert, O. (1998) The taxonomy of developmental control in *Caenorhabditis elegans*. *Science* **282**, 2033-2041.
226. Krishna, S., Maduzia, L. L., and Padgett, R. W. (1998) Specificity of TGF- $\beta$  signaling is conferred by distinct type I receptors and their associated Smad proteins in *Caenorhabditis elegans*. *Development* **126**, 251-260.
227. Reddy, K. B., Karode, M. C., Harmony, A. K., and Howe, P. H. (1996) Interaction of transforming growth factor- $\beta$  receptors with apolipoprotein J/clusterin. *Biochemistry* **35**, 309-314.

STIC-LL

N012/5

From: Holleran, Anne  
S nt: Friday, December 05, 2003 1:39 PM  
To: STIC-ILL  
Subject: refs. for 09/780,953

#74436

Please send copies of the following papers:

1. Gold Molecular Neurobiology (1997) 15(3): 285-306
2. Jain Transplant Intl. (2002) 15(12): 630-634
3. Piek Faseb J. (1999) 13(15): 2105-2124

12478410

Anne Holleran  
AU: 1642  
Tel: 308-8892  
RM: 8e03

mailbox: 8e12

Sunjay Jain  
Mostafa A. S. Mohamed  
Rebecca Sandford  
Peter N. Furness  
Michael L. Nicholson  
David Talbot

## Sequential protocol biopsies from renal transplant recipients show an increasing expression of active TGF $\beta$

Received: 8 October 2001  
Revised: 5 June 2002  
Accepted: 9 July 2002  
Published online: 19 October 2002  
© Springer-Verlag 2002

S. Jain (✉) · R. Sandford · P.N. Furness  
M.L. Nicholson  
University Department of Surgery,  
Leicester General Hospital, Gwendolen  
Road, Leicester, LE5 4PW, UK  
E-mail: sj34@le.ac.uk  
Tel.: +44-116-258 8080  
Fax: +44-116-249 0064

M.A.S. Mohamed · D. Talbot  
Department of Surgery, University  
of Newcastle, Freeman Hospital,  
Newcastle-upon-Tyne, NE7 7DN, UK

**Abstract** Chronic allograft nephropathy (CAN) is a major cause of graft loss after renal transplantation. Implicated in the pathogenesis of this complication is overproduction of the cytokine transforming growth factor beta (TGF  $\beta$ ). In this study we measured changes in CAN's expression in stable patients early after transplantation, and studied links with established risk factors for CAN, such as delayed graft function, acute rejection, and cyclosporine exposure. We took biopsies from 40 renal allografts at time of transplantation (pre-perfusion), and then, using ultrasound guidance, at 1 week and 6 months after transplantation. An immunofluorescence technique was used to stain sections for active TGF  $\beta$ . These were then assessed by semi-quantitative scanning laser confocal microscopy. There was very little variation in active TGF- $\beta$  expression among patients in their pre-perfusion biopsies.

Expression had increased by 1 week and then very significantly by 6 months ( $P < 0.0001$ ). Patients who suffered delayed graft function had increased TGF- $\beta$  expression at both time points. There was no difference regarding donor type, acute rejection, and immunosuppressive drug (cyclosporine or tacrolimus). There was no correlation between the amount of TGF- $\beta$  expression at any time-point and isotope glomerular filtration rate (GFR) at 12 months. This study demonstrated that in a group of stable renal allograft recipients, TGF- $\beta$  expression in the kidney increased after transplantation. As the study used protocol biopsies, this increase is unlikely to be due to acute events, and probably represents a genuine increase.

**Keywords** Chronic allograft nephropathy · Transforming growth factor beta · Protocol biopsies

### Introduction

As the use of modern immunosuppressives has resulted in less early graft loss due to acute rejection, many renal transplants succumb to a gradual deterioration several years later. This is characterised histologically by fibrosis affecting all compartments of the kidney, and is most commonly described as chronic allograft nephropathy (CAN). While the aetiology is incompletely understood, evidence from other forms of renal fibrosis suggests that

a crucial common pathway is a disturbance in the control of extracellular matrix (ECM) turnover. While this control is complex, one cytokine, transforming growth factor beta (TGF  $\beta$ ), has been consistently shown to play a vital role.

TGF  $\beta$  has been studied extensively in experimental models of renal disease and increased expression and shown to be associated with histological changes of fibrosis [22]. In transplantation, there is much circumstantial evidence to support its role in CAN [8], however,

much about its overall significance in the process remains unanswered. One of the reasons for this is that it does play an immunosuppressive role, and, therefore, some of the up-regulation after transplantation may be beneficial [12].

As yet there have been no reports examining TGF- $\beta$  expression in the same kidney sequentially from before transplantation and further into the early post-transplant period. This is of some importance, because it is likely that genotypic differences mean that individual recipients have different baseline levels that are related to their susceptibility to fibrosis [1]. Comparison between patients may, therefore, be unreliable. A greater understanding of the natural history of TGF- $\beta$  expression after renal transplantation would put into a better context the changes that are found during disease processes such as CAN.

We previously studied the expression of active TGF  $\beta$  after renal transplantation using an immunofluorescence technique [14]. The aim of the present study is to quantify the changes in TGF- $\beta$  expression after transplantation using protocol biopsies at pre-perfusion, at 1 week and at 6 months after transplantation.

### Patients and methods

A consecutive series of 40 patients undergoing renal transplantation underwent protocol biopsies. Baseline biopsies (pre-perfusion) were taken from each kidney at the time of transplantation, and subsequently tru-cut biopsies were taken using ultrasound guidance at 1 week and at 6 months after transplantation. Patients whose grafts had failed prior to a 6-month protocol biopsy were not included in the study. All patients underwent regular clinic follow-up, and renal function was quantified by use of the isotope glomerular filtration rate (GFR) measured by  $^{51}\text{Cr}$ -EDTA at 6 and 12 months after transplantation. For immunosuppression we implemented a dual therapy with calcineurin inhibitor (cyclosporine or tacrolimus) and prednisolone. Patients with kidneys from non-heart-beating donors received reduced doses of calcineurin inhibitors and azathioprine (triple therapy). No patients were administered angiotensin converting enzyme (ACE) inhibitors or angiotensin II receptor antagonists. Biopsy specimens were all assessed by an experienced histopathologist (PNF) and graded according to Banff criteria [16].

### TGF- $\beta$ immunofluorescence staining

Paraffin-processed tissue sections were de-waxed and re-hydrated before being incubated with 'blocking' serum (normal human serum 1/15 and normal rabbit serum 1/5 in Tris-buffered saline (TBS) pH 7.6) for 2 h at 4 °C. This was followed by incubation with the primary antibody, chicken anti-human TGF  $\beta$ 1 (RD systems) at 1/100 in blocking serum, overnight at 4 °C. The specificity of this primary antibody had previously been shown by its capacity to block the biological activity of TGF  $\beta$  in an epithelial-cell proliferation assay [13]. After having been washed in TBS for 10 min, sections were incubated for 2 h at 37 °C with rabbit anti-chicken IgG conjugated to fluorescein isothiocyanate at 1/400 in blocking serum. Sections were then washed for a further 10 min in TBS and mounted in fluorescence mounting medium (DAKO, Ely, Cambs.,

UK). Matched negative controls were prepared by replacement of the primary antibody with normal anti-chicken IgG.

Sections were analysed by semi-quantitative scanning laser confocal microscopy as described previously [17]. Data were expressed as the ratio of mean fluorescence over the selected area of experimentally stained tissue (excluding the tubule lumen) to the corresponding value in control sections.

### Statistics

Values of TGF- $\beta$  staining intensity over time were compared using the Wilcoxon signed rank test. Comparisons between different groups were made with the Kruskal-Wallis test for comparison of non-parametrically distributed variables.

### Results

Details of the patients studied are shown in Table 1; those of the histological findings in the protocol biopsies in Table 2. Histological results were mostly normal, however occasional sub-clinical rejection was found. Over 25% of the 6-month biopsies showed evidence of fibrosis.

The most striking finding was the highly significant increase in active TGF- $\beta$  staining 6 months after transplantation (Fig. 1). When the levels of the individual patients were plotted, 34 showed an increase in TGF  $\beta$  over time and only six a drop (data not shown). The TGF- $\beta$  levels both at pre-perfusion and 1 week after transplantation were similar and did not show a large range. TGF- $\beta$  levels did not seem to be affected by donor type (Fig. 2). Although there was only a small number of patients in this study, an attempt was made to determine whether any of the established prognostic factors for graft outcome were related to the rise in TGF  $\beta$ . Only delayed graft function (DGF) seemed to be relevant; patients with DGF displayed increased TGF- $\beta$  expression at both 1 week and at 6 months, when compared with those who did not, however, neither of these results quite reached significance (Fig. 3, 4). Six patients developed acute rejection in the first 6 months after transplantation; they did not have higher levels of TGF  $\beta$  at 6 months (Fig. 5). The type of immunosuppressive drug administered did not affect 6-month

**Table 1** Patient details ( $n=40$ ) (CAD cadaveric donor, NIBD non-heart-beating donor, LRD living related donor, CIT cold ischaemia time)

Characteristic	
Mean donor age in years	42.5 (14.2)
Mean recipient age in years	44 (12.7)
Gender ratio (M:F)	24:16
Donor type (CAD; NIBD; LRD)	17; 14; 9
Mean CIT	12.3 (8.0)
DGF	14/40 (35%)
AR by 6 months	6/40 (15%)

**Table 2** Histological findings in protocol biopsies (according to the Banff 1997 classification)


Diagnosis	Biopsy		
	Pre-perfusion	1 week	6 months
Normal	36	21	26
Acute tubule necrosis	0	13	0
Borderline rejection	0	4	3
AR	0	2	0
CAN	4	0	11

TGF- $\beta$  levels either (Fig. 6), and there was no correlation with drug levels for either agent (data not shown). There was no correlation between TGF- $\beta$  levels at any time-point and GFR at 6 or 12 months after transplantation.

### Discussion

This study demonstrates increased levels of active TGF  $\beta$  in renal transplants at 6 months when compared with baseline pre-perfusion levels. As the study was carried out using protocol biopsies, changes were likely to be due to transplantation rather than to any particular condition for which the biopsies had been taken. This study comprises a relatively small number of patients and so does not have the power to demonstrate differences between patient groups after transplantation. Furthermore, it does not show any influence of TGF- $\beta$  levels on prognosis. This will need larger studies.


There have been problems with the study of TGF- $\beta$  expression in biopsy specimens because of uncertainties about whether staining distinguished between active and latent forms. The antibody used in this study has been shown to react only with active TGF  $\beta$ ; this study was therefore carried out optimally [13].

**Fig. 1** Changes in TGF- $\beta$  expression after transplantation

Although several human studies have reported increases in the expression of TGF  $\beta$  in CAN [15, 18, 19], none of these drew comparisons with stable post-transplant patients. Hence, it is not clear whether TGF  $\beta$  is truly involved in the process, or if it is an innocent bystander.

There have been few studies of renal TGF- $\beta$  expression after transplantation in stable recipients. Lantz et al. included the biopsies of seven stable transplant recipients 6–24 months after transplantation in their series of 28 biopsies stained for TGF  $\beta$  [10]. Those showed definitely increased staining when compared with normal controls, measured in a semi-quantitative manner on a scale of + to + + +. No difference was shown between stable transplants and those with acute rejection (AR) or CAN. The only other study of TGF- $\beta$  expression in stable patients after transplantation measured its levels in plasma. [5] There was a definite increase in the 17 transplant patients when compared with 43 healthy controls ( $P=0.0004$ ). Again, no difference was determined between the stable transplants and those with AR or CAN. These findings are in agreement with those in the current study, which shows that while there was a definite increase in TGF  $\beta$  at 6 months, levels were not affected by established risk factors for CAN.

There is a very good reason for increased TGF- $\beta$  levels in post-transplant patients, as it has an immunomodulatory role and is likely to be significant for reducing the extent of the reaction against the allograft [12]. It also seems to be up-regulated by ischaemia and to have a protective role [11]; indeed this may explain the increased levels found in patients with DGF in this series. As well as potential beneficial actions however, TGF  $\beta$  does play a role in fibrosis, which has been described as its dark side [2]. Possibly, certain patients are more prone to the unwanted effects of TGF  $\beta$ , and an interesting finding with regard to this has been that

**Fig. 2** Active TGF- $\beta$  expression 6 months after transplantation, according to donor type






Fig. 3 Effect of DGF on active TGF- $\beta$  expression 1 week after transplantation




Fig. 4 Effect of DGF on active TGF- $\beta$  expression 6 months after transplantation




Fig. 5 Effect of AR on active TGF- $\beta$  expression 6 months after transplantation




Fig. 6 Active TGF- $\beta$  expression 6 months after transplantation, according to primary immunosuppressant drug

there are various genotypes of TGF  $\beta$ , with differing propensities for developing fibrosis [1].

The calcineurin inhibitors cyclosporine and tacrolimus induce TGF  $\beta$  [20, 21], and this has been proposed as a mechanism of action [9]. There has been much interest in differences between the two drugs, and indeed, we have previously shown that cyclosporine induced TGF  $\beta$  to a greater extent in biopsies taken for diagnostic purposes. A study using protocol biopsies after liver transplantation also showed this, and thus it is surprising that there was no difference found in the current study. However, all these studies comprise relatively small numbers, and so they are probably not conclusive. There have certainly been other studies that could not find a difference in the effects of the two drugs on TGF  $\beta$ , either [7]. ACE inhibitors have been shown to have a down-regulatory effect on TGF- $\beta$  expression [3]. While there were no patients on these drugs in this

study, they may have had a confounding effect when drug groups in other studies were compared. The use of angiotensin II receptor blockers therapeutically to reduce TGF- $\beta$  levels after renal transplantation is an interesting concept that is being explored [4].

A correlation between increased TGF- $\beta$  levels and rate of decline in renal function has been shown by Cuhaci et al. [6]. In this study of forty patients, TGF- $\beta$  levels were graded as low or high in a qualitative manner. There was no correlation between TGF- $\beta$  levels at 1 week and at 6 months and renal function at 6 months and at 1 year in this study. Obviously longer follow-up will allow a more detailed analysis to be made.

In conclusion, there still remains much to be learned about the importance of TGF  $\beta$  after renal transplantation, and its role needs to be defined more clearly before its manipulation for potential therapeutic gain is a practical option.

## References

1. Awad MR, El-Gamel A, Hasleton P, Turner DM, Sinnott PJ, Hutchinson IV (1998) Genotypic variation in the transforming growth factor-beta1 gene: association with transforming growth factor-beta1 production, fibrotic lung disease, and graft fibrosis after lung transplantation. *Transplantation* 66:1014-1020
2. Border WA, Ruoslahti E (1992) Transforming growth factor-beta in disease: the dark side of tissue repair. *J Clin Invest* 90:1-7
3. Border WA, Noble NA (1998) Interactions of transforming growth factor-beta and angiotensin II in renal fibrosis. *Hypertension* 31:181-188
4. Campistol JM, Inigo P, Jimenez W, Lario S, Ciesca PH, Oppenheimer F, Rivera F (1999) Losartan decreases plasma levels of TGF-beta1 in transplant patients with chronic allograft nephropathy. *Kidney Int* 56:714-719
5. Coupes BM, Newstead CG, Short CD, Brenchley PE (1994) Transforming growth factor beta 1 in renal allograft recipients. *Transplantation* 57:1727-1731
6. Cuhaci B, Kumar MS, Bloom RD, Pratt B, Haussman G, Laskow DA, Alidoost M, Grotkowski C, Cahill K, Butani L, Sturgill BC, Pankewycz OG (1999) Transforming growth factor-beta levels in human allograft chronic fibrosis correlate with rate of decline in renal function. *Transplantation* 68:785-790
7. Hughes JR, Hughes VF, Trull AK, Metcalfe SM (1999) Blood levels of TGF beta1 in liver transplant recipients receiving either tacrolimus or microemulsified cyclosporine. *Transplantation* 68:583-586
8. Jain S, Furness PN, Nicholson ML (2000) The role of transforming growth factor beta in chronic allograft nephropathy. *Transplantation* 69:1759-1766
9. Khanna A, Li B, Sharma VK, Suthanthiran M (1996) Immunoregulatory and fibrogenic activities of cyclosporine: a unifying hypothesis based on transforming growth factor-beta expression. *Transplant Proc* 28:2015-2018
10. Lantz I, Dimeny E, Larsson E, Fellstrom B, Funa K (1996) Increased immunoreactivity of transforming growth factor-beta in human kidney transplants. *Transpl Immunol* 4:209-214
11. Lefer AM (1991) Mechanisms of the protective effects of transforming growth factor-beta in reperfusion injury. *Biochem Pharmacol* 42:1323-1327
12. Letterio JJ, Roberts AB (1998) Regulation of immune responses by TGF-beta. *Annu Rev Immunol* 16:137-161
13. Mohamed MAS, Walmsley M, Robertson H, Kirby JA, Talbot D (1999) The effect of cyclosporin A and tacrolimus on cultured human epithelial cells: the role of TGF beta. *Transplant Proc* 31:1173
14. Mohamed MAS, Robertson H, Booth TA, Balpuri S, Kirby JA, Talbot D (2000) TGF beta expression in renal transplant biopsies. A comparative study between cyclosporin A and tacrolimus. *Transplantation* 69:1002-1005
15. Pankewycz OG, Miao L, Isaacs R, Guan J, Pruett T, Haussmann G, Sturgill BC (1996) Increased renal tubular expression of transforming growth factor beta in human allografts correlates with cyclosporine toxicity. *Kidney Int* 50:1634-1640
16. Racusen LC, Solez K, Colvin RB, Bonsib SM, Castro MC, Cavallo T, Croker BP, Demetris AJ, Drachenberg CB, Fogo AB, Furness P, Guber LW, Gibson IW, Glotz D, Goldberg JC, Grande J, Halloran PF, Hansen HE, Hartley B, Hayry P, Hill CM, Hoffman EO, Hunsicker LG, Lindblad AS, Marcussen N, Mihatsch MJ, Nadasdy T, Nickerson P, Olsen TS, Papadimitriou JC, Rhandhawa PS, Rayner DC, Roberts I, Rose S, Rush D, Salinas-Madrigal L, Salomon DR, Sund S, Taskinen E, Trpkov K, Yamaguchi Y (1999) The Banff 97 working classification of renal allograft pathology. *Kidney Int* 55:713-723
17. Robertson H, Morley AR, Talbot D, Callanan K, Kirby JA (2000) Renal allograft rejection: beta chemokine involvement in the development of tubulitis. *Transplantation* 69:684-687
18. Sharma VK, Bologa RM, Xu GP, Li BG, Mouradian J, Wang J, Serur D, Rao V, Suthanthiran M (1996) Intra-graft TGF-beta 1 mRNA: a correlate of interstitial fibrosis and chronic allograft nephropathy. *Kidney Int* 49:1297-1303
19. Shihab FS, Yamamoto T, Nast CC, Cohen AH, Noble NA, Gold LI, Border WA (1995) Transforming growth factor-beta and matrix protein expression in acute and chronic rejection of human renal allografts. *J Am Soc Nephrol* 6:286-294
20. Shihab FS, Bennett WM, Tanner AM, Andoh TF (1997) Mechanism of fibrosis in experimental tacrolimus nephrotoxicity. *Transplantation* 64:1829-1837
21. Shin GT, Khanna A, Ding R, Sharma VK, Lagman M, Li BG, Suthanthiran M (1998) In vivo expression of transforming growth factor-beta1 in humans: stimulation by cyclosporine. *Transplantation* 65:313-318
22. Yamamoto T, Noble NA, Miller DE, Border WA (1994) Sustained expression of TGF-beta1 underlies development of progressive kidney fibrosis. *Kidney Int* 45:916-927